

18-1964; 1 ♀, Grassland, 750 m., V-5-1964 (Cavagnaro). FLOREANA: 1 ♂, 1 ♀, II-18-1964 (Usinger); 1 ♀, Wittmer Farm, II-15-1964 (Usinger).

This species differs from *P. insularis* by the color of body, and by the number and location of spots on the posterior femora.

### 7. *Psallus insularis* Barber.

(Figures 9a, 14.)

*Psallus insularis* BARBER, 1925, Zoologica, vol. 5, p. 250.

Characterized by its color and number of black spots on posterior femora.

MALE. Length 2.5 mm., width 1.1 mm. Head: Length 0.1 mm., width 0.6 mm., vertex 0.30 mm. Antennae: Segment I, length 0.1 mm.; II, 1.0 mm.; III, 0.5 mm.; IV, 0.3 mm. Pronotum: Length 0.4 mm., width at base 0.9 mm.

General color sordid stramineous to pale yellow; eyes dark brown; posterior femora with four to five larger black spots on inferior margin and a few smaller ones on external surface; tibial spines and spots at their bases, black (fig. 8a). Body covered by black bristles and silvery scale-like hairs.

Rostrum reaching to hind coxae.

Genitalia: Aedeagus (fig. 14a) with vesica as illustrated. Left clasper (fig. 14b) and right clasper (fig. 14c) as shown in figures.




FIGURE 14. *Psallus insularis* Barber: a, Aedeagus; b, Left clasper; c, Right clasper.