

ranges): Length, 3.60 mm (3.67 mm, 3.52-3.90 mm); width, 1.60 mm (1.55 mm, 1.50-1.64 mm). Head length, 0.32 mm (0.33 mm, 0.28-0.38 mm); head width through eyes, 0.84 mm (0.82 mm, 0.78-0.86 mm); vertex width, 0.50 mm (0.48 mm, 0.46-0.50 mm). Length of antennal segment I, 0.28 mm (0.27 mm, 0.26-0.28 mm); II, 1.16 mm (1.08 mm, 1.02-1.16 mm); III, 0.76 mm (0.71 mm, 0.70-0.76 mm); IV, 0.38 mm, 0.36 mm, 0.34-0.38 mm). Pronotal length, 0.60 mm (0.55 mm, 0.54-0.60 mm); width, 1.26 mm (1.21 mm, 1.18-1.26 mm). Cuneal length, 0.70 mm (0.65 mm, 0.64-0.70 mm); width, 0.62 mm (0.59 mm, 0.56-0.62 mm).

Similar to male in form and color.

HOLOTYPE: male, Nova Teutonia, Santa Catarina, Brazil, 27°11'N, 52°23'W, December, 1970, Fritz Plaumann. Deposited in the *J. C. M. Carvalho collection*, Rio de Janeiro.

RO. ALLOTYPE: female, same data as holotype. Deposited in the *J. C. M. Carvalho collection*.
PARATYPES: 5 males, 6 females, same locality; collected September, October, November, December, 1970 and July, November, 1971. Deposited in the collections of the authors.


This species exhibits variation in color. Some specimens are greenish while others are pale yellow. The dark brown scutellum and dark fuscous membrane are common to all individuals.

Slaterocoris reinhardi n. sp.

(Figs. 27-30)

Characterized by its color, dimensions and structure of the male genitalia.

MALE (measurements taken from 5 specimens; those of holotype given first



Slaterocoris reinhardi n. sp. Fig. 27 — male, paratype; fig. 28 — penis; fig. 29 — left paramere; fig. 30 — right paramere.