

Rhinomiris vicarius (Walker, 1873)
Kirkaldy, 1902

One specimen, Carin Chebá. Larger than *camelus* Poppius, calli with only median line pale, pronotum with three longitudinal lines or fasciae reaching the calli.

Trichofulvius fasciatus Poppius, 1909

Two specimens, New Guinea. Brown with a median transversal pale fascia running along middle of clavus, corium and embolium, a spot on corial commissure and a large spot on apex of exocorium and embolium ochraceous. The species belongs to genus *Halldapus* Fieber.

MUSEUM OF NATURAL HISTORY, WIEN

Aspidobothrus dimidiatus (Stål, 1860)
Reuter, 1907

Brasilien, Schott, 2 females. Head, pronotum and mesoscutum reddish lutescent; coxae and base of femora pale.

Aspidobothrus grandis Reuter, 1907

Rio Grande do Sul, 1 male, 1 female, Schott, Brasilien. Head, pronotum, mesoscutum, extreme base of hemelytra reddish lutescent; femora II pale at base, mesosternum with a lateral lutescent spot (hind femora broken). This species was synonymized with *basalis* (Walker, 1873) by Carvalho, 1954: 424.

Aspidobothrus latipennis Reuter, 1907

One specimen, female, Stieglmayr, Rio Grande do Sul. Head and pronotum reddish lutescent, side of mesosternum and anterior coxae reddish.

Aspidobothrus robustus Reuter, 1907

One specimen, female, Brasilien. Large, with a black spot median spot behind calli, mesosternum lutescent to reddish laterally, first coxa red. As mentioned previously (Carvalho, 1954: 424) it is a synonym of *designatus* (Distant, 1888) described in the genus *Resthenia* Spinola.

Aspidobothrus ruficeps (Berg, 1878)
Reuter, 1909

Plata, Berg, 1 male. A large black spot behind middle of calli, mesoscutum, extreme base of hemelytra lutescent, as well as, head and pronotum; sides of mesosternum lutescent.

Aspidobothrus signaticollis Reuter, 1907

Rio Grande do Sul, Stieglmayr, 1 male. With a black spot behind middle of calli; extreme base of hemelytra lutescent; head and pronotum lutescent to reddish, sides of propleura and mesosternum reddish.

Cariojanus pulchellus (Reuter, 1907)
Carvalho, 1954

One specimen, Hetscheko, Blumenau, '80. Pronotum strongly constricted at middle, narrowed anteriorly, collar wide, segment I of antenna incrassate, segment II slender beset with long hairs; a reddish spot present at middle of corium, apex of embolium, internal margin of cuneus at middle; membrane fuscous at base; eyes contiguous to collar; hind coxae, abdomen at base, middle and apex reddish; rostrum reaching middle coxae, hind femora mutilated.

Caulotops cyaneipennis Reuter, 1908

One specimen, Bilimek, Saint Marcos. Brown to cinnamon, scutellum and head lutescent, legs pale with fuscous spots on anterior femora, sides of frons and clypeus black, segment I of antenna black (remaining segments mutilated); rostrum reaching the hind coxae.

Caulotops puncticollis Bergroth, 1898

Buenos Aires, 1 specimen. Head and calli lutescent, body brown (Col. Signoret).

Cyclidolon cerinum Reuter, 1908

One specimen, Bilimek, Mexico, 1871. Color pale (cerinum), legs pale yellow, second antennal black apically; size small, cuneal fracture large and deep, cuneus strongly rounded externally. The genus has the facies of *Lundiella* Carvalho, 1951.