## Orthocephalus tenuicornis var. fulvipes Reuter 1904

Tarsus (Reuter 1904 b). Known only from the type locality mentioned.

#### Orthocephalus parvulas Reuter 1891

Izmir, Bursa (Reuter 1891 a).

Up till now recorded from the two above-mentioned localities and seems to be a species endemic in the Aegean part of Anatolia.

# Orthocephalus bivittatus Fieber 1864

1 d — Edirne, 8.—13. VI. 1947. (Exp. of the National Museum, Praha.) On vegetation of steppe formation.

This specimen constitutes a new record for Turkey.

Species of Holomediterranean distribution, in SW Asia recorded from Caucasia and Transcaucasia and in C. Asia from Turkestan.

## Orthocephalus vittipennis (Herrich-Schäffer 1835)

4 of and 15 99 — Gerede, 22. VI. 1947. (Exp. of the National Museum, Praha.)

On Anthemideae in the border zone of forest.

Species recorded in Turkey only from Merdenik (Kiritshenko 1918).

Species of Eurosiberian distribution, in SW Asia recorded from Transcaucasia.

## Pachytomella phoenicea (Horváth 1884)

Tarsus (Reuter 1904 b).

Species listed only from Syria, Israel and above-mentioned locality in S. Anatolia, probably species of Pontomediterranean distribution.

# Pachytomella passerinii (Costa 1841)

Bursa (Horváth 1883 a, Reuter 1891 a).

Species of Holomediterranean distribution, in the Middle East recorded only from Israel.

#### Piezocranum corvinum Puton 1895

Gaziantep (Reuter 1895).

Species known only from this type locality. Probably endemic of northern part of the Syrian plateau.