

**Conostethus roseus** (F a l l e n 1829)

4 ♂♂ and 1 ♀ — Toros: Bürücek, 29.—31. VII. 1947. (Exp. of the National Museum, Praha.)

On xerophilous vegetation (*Trifolium* sp.) of wooded hill formation.

Representation of this species in Turkish fauna listed only from Asia Minor (O s h a n i n 1909, 1912).

Species of Holomediterranean distribution, in the Middle East recorded further only from Israel.

**Conostethus venustus** (F i e b e r 1858)

Izmir (R e u t e r 1879 b).

Species of Holomediterranean distribution, in the Middle East recorded from Syria and Cyprus.

**Pronototropis longicornis** R e u t e r 1900

Ekbaz (H o r v á t h 1901).

Species of Holomediterranean distribution, listed from Morocco, Cyprus and Israel.

**Pachyxyphus lineellus** (M u l s a n t a n d R e y 1852)

2 ♀♀ — Toros: Bürücek, 29.—31. VII. 1947. (Exp. of the National Museum, Praha.)

On the xerophilous vegetation of wooded hill formation.

Up till now recorded in Turkey from Bursa (H o r v á t h 1883 a) Belgrad ormani (F a h r i n g e r 1922); Asia Minor (R e u t e r 1789 b). Occurring on *Cistus laurifolius* (F a h r i n g e r).

Species of Holomediterranean distribution, in the Middle East recorded from Israel and Caucasia.

**Thermocoris munieri** P u t o n 1875

1 ♂ — Toros: Bürücek, 29.—31. VII. 1947. (Exp. of the National Museum, Praha.)

On the xerophilous vegetation of wooded hill formation.

Up till now this species has been recorded in Turkey from Bursa (H o r v á t h 1883 a, R e u t e r 1878, 1890 a) and Sapanca—Eskişehir (H o r v á t h 1905 a).

Species of Holomediterranean distribution, in the Middle East recorded from Cyprus.

**Opisthotaenia fulvipes** R e u t e r 1901

1 ♀ — Edirne, 8.—13. VI. 1947. (Exp. of the National Museum, Praha.) Collected on the vegetation of steppe formation.

In Turkey so far listed only from Amasya (R e u t e r 1901 a).

Species of Pontomediterranean distribution, in SW Asia recorded from Transcaucasia.