

***Psallus (Nanopsallus) carduellus* var. *quadrisignatus* Reuter 1904**

Yamanlar dağı near Izmir (Reuter 1904 b).

Variety recorded only from the above-mentioned type locality. Typical form is of Pontomediterranean distribution.

***Psallus (Nanopsallus) carduellus* var. *infuscatus* Reuter 1904**

Yamanlar dağı near Izmir (Reuter 1904 b).

Variety recorded only from the above-mentioned type locality.

***Excentricus punctipes* (Fieber 1864)**

Bursa (Horváth 1883 a, Reuter 1883).

Species of Caspian distribution; also recorded from South Russian steppe region.

***Excentricus oophorus* Horváth 1888**

Bursa (Horváth 1888); Asia Minor (Reuter 1913).

Species only known from the above mentioned locality.

***Criocoris sulcicornis* (Kirschbaum 1855)**

3 ♂♂ and 6 ♀♀ — Moğan gölü, 8. VII. 1947. (Exp. of the National Museum, Praha.)

On *Rubiaceae* in steppe formation.

Up till now recorded in Turkey only from Yilanli dağı near Kayseri (Horváth 1905 a).

Species of Holomediterranean distribution, in SW Asia recorded from Caucasia and Transcaucasia.

***Plagiognathus (Plagiognathus) alpinus* Reuter 1875**

3 ♀♀ — Moğan gölü, 5. VII. 1947. (Exp. of the National Museum, Praha.)

On salt steppe formation.

New record for Turkey.

Species of European distribution, in SW Asia recorded from Caucasia, Transcaucasia, in Central Asia from Turkestan.

***Plagiognathus (Plagiognathus) bipunctatus* Reuter 1883**

1 ♀ — Moğan gölü, 9. VII. 1947. 1 ♂ — Adana, 1.—3. VIII. 1947. 3 ♂♂ — Karataş, 2.—5. VIII. 1947. (Exp. of the National Museum, Praha.)

Up till now recorded in Turkey only from Zardanes (Kiritschenko 1918).