

Collected on *Verbascum* sp. of salt steppe formation (Moğan gölü) and at dusk flying high above the ground in cultivated area and steppe formation.

Species of Holomediterranean distribution, extending in steppe and leafy wood formations into Central Europe; in the Middle East recorded from Cyprus, Caucasia, Transcaucasia, in Central Asia from Tajikistan.

Plagiognathus (Plagiognathus) fulvipennis (Kirschbaum 1855)

4 ♂♂ and 4 ♀♀ — Edirne, 8.—13. VI. 1947. 1 ♀ — Beynam, 28. VI. 1947. 2 ♀♀ — Ankara-Baraj, 3.—4. VII. 1947. 1 ♀ — Gyaur dağlari, 17. VIII. 1947. (Exp. of the National Museum, Praha.)

On *Echium* sp. of steppe and wooded hill formations.

Up till now recorded in Turkey only from Zardanes (Kiritschenko 1918).

Species of Holomediterranean distribution, in SW Asia recorded from Caucasia and Transcaucasia.

Plagiognathus (Plagiognathus) arbustorum (Fabricius 1794)

7 ♂♂ and 5 ♀♀ — Mollafeneri, 21. VI. 1947. (Exp. of the National Museum, Praha.)

On low undergrowth in macchia.

Up till now recorded in Turkey only from Belgrad ormani (Fahring 1922), occurring on *Urtica dioecia*.

Species of Eurosiberian distribution, in SW Asia recorded from Caucasia and Transcaucasia, in Central Asia from Tajikistan.

Utopnia torquata (Putton 1881)

Bursa (Horváth 1883 a, Reuter 1883), Iskenderon (Horváth 1901), Tarsus (Reuter 1904 b), Gyaur dağlari (Horváth 1919), Kusçular, Haruniye (Fahring 1922); Anatolia (Reuter 1890 a).

Species of Pontomediterranean distribution, in the Middle East recorded from Syria and Israel.

Chlamydatus (Attus) pullus (Reuter 1870)

4 ♂♂ and 1 ♀ — Mollafeneri, 21. VI. 1947. (Exp. of the National Museum, Praha.)

Beneath low vegetation in macchia formation.

Up till now recorded in Turkey only from Merdenik (Kiritschenko 1918).

Species of Eurosiberian distribution, in SW Asia recorded from Caucasia, Transcaucasia, in Central Asia from Turkestan.