

Miris Fabr.*M. dolabratus* (L.)

Cimex dolabratus L. Syst. Nat. ed. 10, 1767, 449. — *Miris d.* v. D. 300. — Bl. 672. — Mo. 20.

W. Glenbournie, Bonne Bay, 19. VIII. 49, 20 ind. (P); Lomond, 14. VII. 49, 3 ind. (K).

E c o l o g y: On meadows; in Glenbournie found on *Dactylis glomerata*. In Lomond on a humid meadow with grasses, *Heracleum*, *Achillea*, *Thalictrum* and *Juncus*.

D i s t r i b u t i o n: Palearctic species, introduced from Europe. From Newfoundland and Quebec to British Columbia, from New England to Minnesota.

Stenodema Lap.*S. trispinosum* Reut.

Stenodema trispinosum Reut. Öfv. Fi. Vet. Soc. Förh. 46, 15, 1904, 4, 8. — v. D. 303. — Bl. 669. — Mo. 20. — Strickl. 201.

Nova Scotia, Baddeck, 24. V. 51, 1 ind. (L); North Sydney, 27. V. 51, 1 ind. (L).

S.E. Holyrood, 11. VI. 49, 4 ind. (K); St. Johns, 5. VI. 49, 3 ind. (K); Waterford Bridge, 5. VI. 49, 5 ind. (P). — S. Rencontre West, 17—19. VI. 49, 1 ind. (L); Cinq Cerf River, 16. VI. 49, 1 ind. (K). — W. St. Fintans, 3. VII. 49, 1 ind. (L); Steady Brook, Humber, 10. VII. 49, 1 ind. (P)? Deer Lake, 1—2. VI. 51, 6 ind. (L); Lomond, 4—5. VI. 51, 1 ind. (L). — N.E. Paquet, 18. VI. 51, 1 ind. (L). — E. Goobies, 15. VIII. 51, 1 ind. (L). — C. Millertown, 10—14. VI. 51, 1 ind. (L).

E c o l o g y: Humid meadows, marshes with *Carex*, *Juncus*, *Typha*, *Iris*, etc. Both on waste places and in woodland.

D i s t r i b u t i o n: Holarctic species. In the northern parts of the palearctic region: Northern Europe (not common), Turkestan, Siberia, Japan. Newfoundland, Quebec to British Columbia, in the northern States from New England to California.

S. vicinum (Prov.)

Miris vicina Prov. Nat. Can. IV, 1872, 77. — *Stenodema v.* v. D. 304. — Bl. 669. — Walley, Can. Ent. LXII, 76. — Mo. 20. — Strickl. 201.

S.E. Holyrood, 11. VI. 49, 4 ind. (K); Portugal Cove, 6. VI. 49, 3 ind. (L); St. Johns, 5. VI. 49, 3 ind. (K); Waterford Bridge, 5. VI. 49, 3 ind. (P). — S. Rencontre West, 16. VI. 49, 7 ind. (L); Terrenceville, 14. VIII. 51, 2 ind. (L). — W. South Branch, 3. VII. 49, 1 ind. (L). — N.E. Paquet, 18. VI. 51, 2 ind. (L); Lewisporte, 26—27. VI. 51, 1 ind. — E. Gambo, 26. VIII. 49, 1 ind. (L); Goobies, 15. VIII. 51, 1 ind. (L). — C. Gander, 2. VI. 49, 2 ind. (P); Millertown, 10—14. VI. 51, 1 ind. (L); Victoria Lake, 11—13. VI. 51, 1 ind. (L).

E c o l o g y: Both on moist meadows with *Carex*, etc., and on dry grassland; on cultivated land in built-up areas.

D i s t r i b u t i o n: Nearctic species with a wide range. From Newfoundland and Quebec to British Columbia, from New England to California. Not recorded from the southernmost States.

Teratocoris Fieb.*T. herbaticus* Uhl.

Teratocoris herbaticus Uhl. Entom. Amer. III, 1887, 67. — v. D. 308. — Bl. 678. — Walley, Can. Ent. LXII, 76. — Mo. 20.

S.E. Labrador: Red Bay, 14—17. VII. 51, 1 ind. (L); Pinware River, 19. VII. 51, 1 ind. (L); L'Asne au Loup, 16. VII. 51, 3 ind. (L).

N.E. Twillingate, 7. VII. 51, 1 ind. (L). — C. Grand Falls, 26. VII. 51, 1 ind. (L).