

*W.* Corner Brook, 16. VIII. 49, 3 ind. (L). — *C.* Millertown Junction, 21. VIII. 49, 14 ind. (K).

*E c o l o g y*: In Corner Brook on *Corylus* and *Sambucus*.

*D i s t r i b u t i o n*: Fairly widespread nearctic species. From Newfoundland, Quebec and New England west to Manitoba, Colorado and Kansas.

*P. c h r y s a n t h e m i* (Wolff)

*Lygaeus chrysanthemii* Wolff, Abb. der Wanz. 1804, fig. 151. — *Plagiognathus c.* Bl. 943. — Mo. 23.

*S.E.* St. Johns, 28. VIII. 49, 1 ind. (L). — *S.* St. Pierre, 11. VIII. 51, 1 ind. — *E.* Terra Nova, 26—28. VII. 51, 1 ind. (L).

*E c o l o g y*: On dry ground influenced by cultivation.

*D i s t r i b u t i o n*: Palearctic species, distributed in Europa, Caucasia, Turkestan and West Siberia. Introduced from Europe to Newfoundland, Quebec, Ontario and New England.

#### *C h l a m y d a t u s* Curt.

*C. pulicarius* (Fall.)

*Lygaeus pulicarius* Fall. Monogr. Cimic. Suec. 1807, 97. — *Chlamydatus p.* v. D. 416. — Bl. 947. — Mo. 24. — Strickl. 199.

*S.E.* St. Johns, 28. VIII. 49, 1 ind. (L). — *N.E.* Norris Arm, 24. VIII. 49, 1 ind. (P); Twillingate, 2—8. VII. 51, 5 ind. (L); Fogo, Fogo, 29. VI. 51, 2 ind. (L).

*E c o l o g y*: On meadows and on margins of cultivated land. Both on dry and humid localities.

*D i s t r i b u t i o n*: Evidently generally distributed in Newfoundland. Quebec, Alberta, New York. Probably introduced palearctic species.

### SALDIDAE

#### *L a m p r a c a n t h i a* Reut.

*L. crassicornis* (Uhl.)

*Salda crassicornis* Uhl. Bull. U. S. Geol. Geogr. Surv. III, 1877, 441. — v. D. 447. — Bl. 1016. — Mo. 26.

*S.E.* Come-by-Chance, 27. VIII. 49, 2 ind. (P). — *N.W.* Port au Choix, 5. VIII. 49, 2 ind. (P); Flowers Cove, 23—25. VII. 49, 2 ind. (L). — *N.E.* Twillingate, 7. VII. 51, 1 ind. (L); Fogo, Fogo, 2. VII. 51, 2 ind. (L); Fogo, Seldom, 3. VII. 51, 1 ind. (L).

*E c o l o g y*: On humid marshes, on swamps with *Carex*, *Myrica*, etc.; on shores of *Sphagnum* pools.

*D i s t r i b u t i o n*: Nearctic species with northern distribution. Newfoundland, Quebec, Saskatchewan; New York, New Hampshire, Nebraska, Indiana.

#### *P e n t a c o r a* Reut.

*P. ligata* (Say)

*Acanthia ligata* Say, Heter. N. Harm. 1832, 34. — *Pentacora* v. D. 439. — Bl. 1005. — Clark, Can. Ent. LVIII, 204. — Walley, Can. Ent. LXII, 77. — Mo. 25.

*E.* Terra Nova, 26—28. VI. 51, 1 ind. (L).

*E c o l o g y*: On sandy ground.

*D i s t r i b u t i o n*: Nearctic species with relatively northern range. From Newfoundland, Quebec and New England west to Manitoba and Minnesota.