Female unknown.

HOLOTYPE: Macropterous &, SOUTH AFRICA: Cape Province, Grootfontein, Middelburg, October, M. Johannsmeier (SANC).

PARATYPES: 3 macropterous 3 3, same data as holotype (SANC, RTS).

This species is named for the collector, M. Johannsmeier.

Coatonocapsus johannsmeieri is similar in coloration to transvaalensis but is much smaller and has the hemelytra shorter relative to the total length of the body. It is much lighter and more elongate than sweeti, and much darker than pallidus.

## Coatonocapsus pallidus, new species

Figures 59, 242

MACROPTEROUS MALE: Basic coloration light yellow green; head with an orangish tinge; all setiform hairs on dorsum with small, round, brown spots at bases; vertex mottled with brown; clypeus, apex of juga and lora, antennae, labium, coxae, mesoscutum, abdomen (basal half greenish), and tarsi brown; femora with many small, round, brown spots; tibiae with dark brown spots at bases of spines.

Labium reaching between procoxae and mesocoxae; posterior margin of pronotum straight.

MALE GENITALIA: Figure 242.

BRACHYPTEROUS FEMALE: Coloration, body surface texture, and vestiture as in male; abdomen yellow green; ovipositor brown. Structurally very similar to female of *C. sweeti*.

FEMALE GENITALIA: See generic discussion.

MEASUREMENTS: Total length 3.08, maximum width 1.32, width head .78, interocular space .40.

HOLOTYPE: Macropterous 3, SOUTH AFRICA: Cape Province, East of Pakhuis Pass, Mus. Expd., Sept. 1947 (SAM).

PARATYPES: Cape Province—1 macropterous  $\delta$ , 4 brachypterous  $\Im$ , 5, 000 Doorn River, XI.1931 (Mackie); 12 brachypterous  $\Im$ , Michells Pass, Ceres Div., Oct. 1934; 1 macropterous  $\delta$ , R. Sonder