

reque longum ac latum. Hemelytra apicem abdominis sat longe superantia, cuneod eclivi, cellula majore membranae apicem cunei paullum superante, apice acutangulo. Abdomen hemelytris æque latum vel paullulo latius. Pedes parce setosi, coxis anticis medium mesosterni vix attingentibus, spinulis tibialibus latitudini tibiarum subaeque longis. — ♂. Long. 4.6, cum membr. 6 mill.

Var. : Pronotum a basi ultra medium et scutellum, maculis duabus basalibus et apice exceptis, nigricantia. Corium totum et clavus, apice excepto, infuscata.

Two or three species of the genus *Adelphocoris* REUT. are known from North America, but no neotropical species has hitherto been recorded. Some of the species described under other generic names by STÅL, BERG and DISTANT may, however, belong here.

Syinias fulvicollis FABR.

The blackish basal spot to the pronotum is sometimes very small and often lacking. The synonymy of this species is :

Capsus fulvicollis FABR., Syst. Rhyng., p. 244.

Ecritotarsus fulvicollis STÅL, Hem. Fabr., I, 85.

Syinias audens DIST., Biol. Cent. Am., Hem. Het. I, 249, tab. XXIV, fig. 25.

The type of FABRICIUS was also from Guiana.

Miccus nov. gen.

Corpus ovatum. Caput fortiter nutans, apice pronoti parum latius, altitudine basali longius, e supero visum late triangulare, altitudine duplo brevius, ab antico visum altitudine nonnihil latius, vertice longitudinaliter impresso et basi carina tenui transversa antrorum levissime curvata marginato, mox ante carinam transversim impresso, clypeo a fronte impressione discreto, basi sat longe infra basin antennarum et paullum infra marginem inferiorem oculorum sita, angulo faciali acutissimo, oculis pronoto contiguis, e latere visis oblique breviter obovatis, genis oculo subæque altis, gula brevi, valde obliqua, rostro crasso, coxas anticis paullum superante, antennis gracilibus, ad oculos nonnihil supra angulum eorum inferiorem insertis, articulo primo apicem clypei paullum superante, secundo primo paullulo longiore, tertio secundo subæque longo, curvato, quarto tertio longiore, gracillimo. Pronotum capite dimidio latius, latitudine sua basali paullulo brevius, lateribus sinuatum, margine apicali subrecto, pone oculos leviter obliquato, area apicali stricturam non formante, sed medio postice inter callos