are to be found around and on the inflated sac like area near terminal opening.

Morphological characters: as given for genus.

Female: identical to male in shape and dimensions.

Holotype: male, Lulua, Kapanga, X-1932, (F. G. OVERLAET coll.). Allotype: female, Elisabethville, II-1940 (H. J. Brédo coll.), in the Collection of the Congo Museum (Tervuren). Paratypes: 5 specimens, Nyangwe, IV-1918 (R. Mayné coll.), Lac Kivu cheff. N'Gwese (Carlier coll.), in the collections of the Congo Museum, British Museum (Nat. History) and the Author.

UELEANA n. gen.

Ortholylinae, Orthotylini. Characterized by the stout first antennal segment coarsely and desenly pubescent and by the very long and somewhat curved cuneus.

Head glabrous, vertex convex, carinate posteriorly, with a short neck; eyes small, rounded, distant from anterior margin of pronotum by a distance equal to about half the width of eye, frons vertical; seen from front the head is triangular, acutely pointed at apex. Rostrum apparently reaching middle coxae (carded specimen).

Antennae inserted contiguous to eye internally, the first joint strongly incrassate about as long as head and pronotum together, strongly narrowed at base, clothed with dense and stout bristles, second segment very slender, linear, shortly and esparsely pubescent, third and fourth mutilated.

Pronotum glabrous, impunctate, faintly rugose transversally, constricted anteriorly, the lateral margins rounded, the posterior margin strongly dented inwards at middle of mesoscutum, calli polished and strongly shining, anterior area imitating a collar, rugose. Mesoscutum partially covered; scutellum flat and small.

Hemielytra short, esparse and semiadpressedly pilose, dilated at middle and constricted at level of cuneal incisure; embolium slightly widened toward apex where it is more or less laminate; cuneus about three times as long as wide, somewhat curved outwards apically; membrane bicellulate, large, with the apex pointed outwards.

Hind legs more developed than the median and anterior pair, the hind femora thickest at middle, tibiae with short spines, tarsi very long and slender, linear, claws slender, arolia convergent at apex, of the Orthotylinae type.

Type of genus: Uealana longicuneata n. sp.