

**Kunungua cinnamomea** n. sp. (Fig. 5, E).

Female : length 3.6 mm., width 1.2 mm. *Head* : length 0.3 mm., width 0.7 mm., vertex 0.40 mm. *Antennae* : segment I, length 0.3 mm.; II, 0.6 mm.; III, 0.7 mm.; IV, 0.7 mm. *Pronotum* : length 0.9 mm., width at base 1.0 mm. *Cuneus* : length 0.5 mm., width at base 0.3 mm.

*Colour* : cinnamon; three last segments of antennae, tarsi and apex of median and posterior tibiae, flavous; tibiae (except apex of middle and anterior pair) black; head, pronotum and base of embolium lighter than hemelytra, the antennal peduncle tending to flavous.

Male : unknown.

*Holotype* : female, Kunungu, 2-IV-1921 (Dr. H. SCHOUTEDEN coll.), in the Collection of the Belgian Congo Museum (Tervuren).

This species differs from *K. boxi* n. sp. by the colour of body, relative length of antennal segments, shorter rostrum not quite reaching anterior coxae, hemelytra widened posteriorly, the tibiae, especially the anterior pair strongly pilose.

**Prodromus melanonotus** n. sp. (Figs. 6 F, A J. K.).

Male : length 4.0 mm., width 1.1 mm. *Head* : length 0.6 mm., width 0.7 mm., vertex 0.34 mm. *Antennae* : segment I, length 0.6 mm.; II, 0.9 mm.; III, 0.9 mm.; IV, 1.7 mm. *Pronotum* : length 0.8 mm., width at base 1.0 mm. *Cuneus* : length 0.8 mm., width at base 0.3 mm.

*Colour* : Hyaline to flavous; head, antennae (except base of first joint), pronotum (except collar), scutellum, piceous; clavus (except the extreme outer basal area), inner margin of corium and embolium, vein of membrane, apex of femora and tibae, fuscous; femora with a dark band on upper surface, apex of head and rostrum yellowish, sternum and abdomen dark to fuscous.

*Genitalia* : aedeagus (fig. 6, F) of the Bryocorinae type. Left clasper (fig. 6 J, K) strongly curved, ended by a point. Right clasper (fig. 6, A) with a few dorsal hairs.

Female : identical to male in colour and dimensions.

*Holotype* : male, Gold Coast, Qm. nr. of Cape Coast, 9 Sept. 1943, H. E. Box coll., H. 1043. *Allotype* : British Cameroons, Mungo River, nr. Tiko, 14 Jan. 1944, H. E. Box coll., in the Collection of the British Museum of Natural History. *Paratype* : male, Haut Uele, Mauda, II-1925, Dr. H. SCHOUTEDEN coll., in the Collection of the Belgian Congo Museum (Tervuren).

This species differs from the others known in the genus by its colour and male genitalia.