

1446. *Capsus pegasus*, sp. n.

Black; basal joint of antennæ, a small discal spot near apex of corium, basal joint of rostrum, coxæ, legs, and disk of abdomen beneath ochraceous; rostrum (excluding basal joint) piceous; tibiæ palely fuscous; pronotum coarsely punctate, with an anterior callosity which is somewhat centrally sulcate; scutellum tumid, transversely foveate at base, obscurely transversely wrinkled; corium coarsely punctate; membrane fuliginous; second joint of antennæ strongly pilose.

Length 5 millim.

Hab. Ceylon (*Lewis*).

1447. *Capsus darsius*, sp. n.

Black; basal joint of antennæ, head beneath, legs, and some transverse fasciæ to abdomen beneath testaceous; second joint of antennæ longly pilose; pronotum strongly punctate, with an obscure anterior callosity which is strongly centrally sulcate; scutellum tumid, transversely foveate at base, punctate and obscurely transversely wrinkled; corium thickly punctate; membrane fuliginous.

Length $4\frac{1}{2}$ millim.

Hab. Burma; Karennee (*Coll. Dist.*).

Genus **BOTHRIOMIRIS**.

Bothriomiris, *Kirk. Tr. Ent. Soc.* 1902, p. 270.

Type, *B. simulans*, *Walk.*

Distribution. Oriental Region.

Oblong, broad, remotely pilose; posterior area of pronotum, scutellum, corium, and sternum thickly strongly punctate; head broad, strongly deflected anteriorly, short as seen above, eyes large, touching the anterior margin of the pronotum but projecting beyond it; antennæ long, second joint longer than pronotum; pronotum without an anterior collar, not distinctly transversely impressed, the anterior area tumid, centrally sulcate, its posterior lateral margins oblique, the posterior area thickly strongly punctate, centrally carinate, the base truncate, oblique towards lateral angles; scutellum somewhat large and subtriangular; cuneus about as broad as long; interior cell of membrane posteriorly angulate; rostrum about reaching the posterior coxæ; femora short, moderately incrassate.

Kirkaldy states that this genus will probably form a new division. I place it here, provisionally, in the *Capsaria*.

1448. *Bothriomiris simulans*, *Walk.* (*Capsus*) *Cat. Het.* vi, p. 125 (1873).

Bothriomiris marmoratus, *Kirk. Tr. E. S.* 1902, p. 271.