joint about twice as long as first; pronotum (excepting the anterior transverse callosity) somewhat coarsely punctate; clavus coarsely, corium less coarsely punctate, costal margin impunctate; membrane longly passing the abdominal apex.

Length 6 mm.

Hab. Mussoorie.

Camptobrochis qualis, sp. n.

Head and pronotum ochraceous, head suffused with sanguineous; eyes black; antennæ with the first and second joints ochraceous, extreme apex of second and the whole of third and fourth joints piceous; pronotum with two transverse callosities near anterior margin and a subbasal linear margin sanguineous; rostrum ochraceous; corium ochraceous, the clavus and apical area of corium sanguineous; cuneus sanguineous, its basal margin pale ochraceous; membrane hyaline, the cellular margins a little darker; body beneath, rostrum, and legs ochraceous; mesosternum and apical areas of femora (more or less) sanguineous; abdomen with a sublateral series of small black spots, in a second specimen these are only faintly visible; antennæ with the first joint a little shorter than head, second nearly twice as long as first; pronotum coarsely punctate and slightly wrinkled, anteriorly deflected; clavus thickly punctate, corium punctate, the outer area almost impunctate; membrane longly passing the abdominal apex; rostrum with its apex black and just passing the intermediate coxæ.

Length 41 mm.

Hab. Simla Hills; Theog, 8000 feet (Annandale).

PACORUS, gen. nov.

Allied to *Camptobrochis*, but with the first joint of the antennæ considerably longer than the head, second joint about twice as long as first and distinctly setose; pronotum twice as broad at base as long and three times as broad as anterior margin; scutellum broad, the basal margin much longer than the lateral margins; corium very broad, excluding cuneus about as long as posterior tibiæ; cuneus with its posterior margin nearly twice as long as its lateral margin; posterior tarsi with the first and second joints subequal in length.

Pacorus politus, sp. n.

Head and pronotum ochraceous, the latter somewhat discoloured on disk and with its posterior margin distinctly

Ann. & Mag. N. Hist. Ser. 8. Vol. iv. 37