

Genus **ASERYMUS**, nov., to replace *Zanessa*. (Vol. II, p. 432.)

Type, *A. sanguinolentus*, Dist.

I originally considered that *Zanessa* Kirk., might include only *sanguinolenta* from Ceylon, but more material has altered my opinion. The description given of *Zanessa* (vol. ii, p. 432) and the figure (vol. ii, fig. 278) will suffice, but stress must be laid on the lateral margins of the pronotum which are obliquely straight and not concavely sinuate as in *Zanessa*, the head is also shorter, broader, and not angularly narrowed anteriorly.

ASERYMUS SANGUINOLENTUS.

Zanessa sanguinolenta, Dist. vol. ii, p. 432, fig. 278.

Genus **DRUTHMARUS.**

(To follow *Zanessa*, Vol. II, p. 432.)

Druthmarus, Dist. *A. M. N. H.* (8) iv, p. 452 (1909).

Type, *D. magnicornis*, Dist.

Distribution. Ceylon.

Head about as long as broad between eyes, centrally longitudinally sulcate, anteriorly broadly angulate; eyes exserted, large, transverse, projecting far beyond the anterior margin of the pronotum, directed a little backwardly; antennæ with the basal joint short and incrassate, second joint very strongly incrassate,




Fig. 133.—*Druthmarus magnicornis*.

about four times as long as first, third and fourth joints short, slender, subequal in length; rostrum reaching the posterior coxæ; pronotum much broader than long, obliquely narrowed to head, moderately deflected anteriorly, and prominently deflected before