

GUNADHYA, gen. nov.

First joint of the antennæ considerably longer than the head, second joint slightly shorter than first and a little more than subequal in length to third joint; pronotum almost twice as broad at base as long and about three times the breadth of anterior margin; scutellum broad and subglobose; corium broad, the lateral margins rounded, ampliate and recurved, cuneus large and ample; the posterior tarsi with the apical joint longest.

Allied to *Pacorus*, Dist., from British India, but with the corium shorter and broader, the lateral margins much more convex, ampliate, and recurved.

Gunadhya rubrofasciata, sp. n.

Reddish ochraceous, lateral margins of corium more sanguineous in hue, cuneus very pale stramineous in hue, its posterior margin very narrowly pale fuscous; antennæ, rostrum, and legs pale stramineous; body beneath imperfectly seen in carded type; pronotum thickly coarsely punctate; scutellum globose, impunctate; lateral margins of corium convex, ampliate, and recurved, with sanguineous reticulations.

Long. 4 mm.

Hab. Upper Houadou R.

Family Pelogonidæ.

Subfam. PELOGONINÆ.

Pelogonus marginatus.

Pelogonus marginatus, Latr. (*Acanthia*) Hist. Ins. xii. p. 242 (1804).

Hab. Gondé; Houadou R.

Fam. Notonectidæ.

Subfam. NOTONECTINÆ.

Enithares bergrothi.

Enithares bergrothi, Montand. Rev. d'Ent. xi. p. 75 (1892).

Hab. Near Dumbea.