

Speciem hanc sub nomine *M. Kiritschenkoi* REUT. in litt. benevole misit nobis clar. A. KIRITSHENKO, in cujus honorem animalculum hoc sub eodem nomine descripsi.

10. *Myrmecophyes lugubris* n. sp.

Brachypterus, elongatus, niger, nitidulus, glaber; capite basi pronoti $\frac{2}{5}$ latiore, vertice plano et quam oculo fere $\frac{2}{3}$ (σ^7) vel fere $\frac{3}{4}$ (φ) latiore; oculis sessilibus; antennis nigris, subglabris, articulo primo latitudine verticis $\frac{4}{5}$ (σ^7) vel dimidio (φ) latiore, lineari, sat gracili, apicem versus setulis nonnullis semierectis obsito, articulo secundo articulo primo $\frac{2^4}{5}$ (σ^7) vel $\frac{2^3}{4}$ (φ) et quam articulo tertio $\frac{3}{4}$ (φ) longiore (articulis quartus deest); rostro nigro-piceo, fere usque ad apicem coxarum posticarum extenso; pronoto transverso, basi longitudine sua $\frac{1}{4}$ (σ^7) vel $\frac{1}{3}$ (φ) latiore, disco convexiusculo, lateribus subrotundatis; mesonoto (sine scutello) pronoto paullo brevior, convexo, a parte basali retrorsum declivi pronoti bene discreto, scutello minuto albo vel flavo-albido, bene discreto; elytris subtiliter punctatis, apice latissime rotundatis et albo-limbatis, limbo albo extrorsum sensim leviter ampliato, angulo apicali interno leviter, externo autem fortius rotundatis, commissura mesonoto $\frac{2}{3}$ — $\frac{3}{4}$ brevior; abdomine nigro-nitente, retrorsum sensim ampliato, in triente apicali (σ^7) vel paullo pone medium (φ) latissimo, marginibus lateralibus rectis (σ^7) vel leviter arcuatis (φ); pedibus nigris, femoribus anticis vel anterioribus interdum apice imo ferrugineis, femoribus posticis margine superiore setis semierectis destituto, tibiis breviter nigro-spinulosis. Long. σ^7 $3^2/3$, φ 4 mill.

Turkestan: Naryn-kol!, leg. G. ALMÁSY 1906. (Mus. Hung.)

Praecedenti valde affinis et simillimus, sed corpore nitidulo, vertice latiore, pronoto minus transverso, mesonoto magis convexo, basi cum basi pronoti haud in eodem plano posito, abdomine nigro-nitente femoribusque posticis margine superiore setis semierectis destituto differt.

11. *Myrmecophyes Korschinskii* REUT.

Jugis et genis feminae basin versus vel totis, interdum etiam clypeo ferrugineo-testaceis; articulo primo antennarum latitudine verticis dimidio (σ^7) vel tantum $\frac{1}{5}$ (φ) longiore, articulo secundo articulo praecedente $\frac{2^4}{5}$ (σ^7) vel $\frac{2^2}{3}$ (φ) et quam articulo tertio $\frac{2}{3}$ (σ^7) vel dimidio (φ) longiore, articulo quarto articulo tertio $\frac{1}{4}$ (φ) brevior; mesonoto (sine scutello) pronoto $\frac{1}{7}$ — $\frac{1}{5}$ brevior, scutello minuto flavo-albido, bene discreto; commissura elytrorum mesonoto dimidio brevior. Long. σ^7 3— $3^1/3$, φ $3^2/5$ —4 mill.