

General color yellow marked with red and brown. Pronotum yellow, calli shining black anteriorly, posterior half with four large brown spots separated by yellow rays; scutellum yellow with two large brown spots each side of median line on apical half; hemelytra somewhat translucent, clouded with red and reddish brown, a distinct brown spot on basal half of corium inside radial vein, apex of corium and embolium with irregular reddish to brown spot, cuneus somewhat translucent, marked with bright red principally on outer basal angle and inner apical angle, membrane dusky, veins pale; sternum with large brown spot; ostiolar peritreme yellow, episternum brown; abdomen with irregular brown spots on sides of segments and with reddish and brown spots on genital segment; legs yellow, tibiae with three reddish to reddish-brown bands, tarsi with apical joint and claws brown, middle and hind femora with broad brown median band. Clothed with fine, short, yellow hairs.

Female. Length 4.9 mm., width 2.3 mm. Head: width 1.2 mm.; vertex .48 mm. Antennae: segment I, length .43 mm.; II, 1.48 mm.; III, .52 mm.; IV, .35 mm. Pronotum: length 1.3 mm., width at base 2.2 mm. Slightly larger and more robust than male. Coloration and pubescence similar to male but color more yellowish, the reddish and brown areas less extensive.

Holotype: male, April 1, 1933, College Station, Texas (H. G. Johnston).

Allotype: Female, taken with the type.

Paratypes: 42 males and females taken with the types on live oak (*Quercus virginiana*) where the species breeds abundantly; 21 males and females, April 24, 1932, Sonora, Texas (S. E. Jones), light trap.

***Neoborus rostratus* n. sp.**

Distinguished by the short antennal segment II which is scarcely equal to width of head through eyes, the long rostrum which slightly surpasses apex of hind coxae, and the lateral carinae of pronotum.

Male. Length 3.5 mm., width 1.57 mm. Head: width .83 mm.; vertex .30 mm.; tylus with black line on apical half which divides on basal half to form two divergent lines that extend upon frons where they become reddish brown, frons shallowly punctate, shining. Rostrum: length 1.48 mm.,