

LEPTOMEROCORIS SIMPLEX.

Leptomerocoris simplex, Walk. Cat. Hem. Het. B. M. vi. p. 145, n. 107 (1873).

Described from Ceylon.

LEPTOMEROCORIS ALBIVIRIDESCENTS.

Leptomerocoris albiviridescens, Motsch. Bull. Mosc. xxxvi. (3) p. 85 (1866).

Described from Ceylon (Patannas).

LEPTOMEROCORIS (?) PISTACINUS.

Leptomerocoris (?) pistacinus, Motsch. Bull. Mosc. xxxvi. (3) p. 85 (1863).

Described from Ceylon (Patannas).

LEPTOMEROCORIS (?) ALBOFASCIATUS.

Leptomerocoris (?) albofasciatus, Motsch. Bull. Mosc. xxxvi. (3) p. 86 (1863).

Described from Ceylon (Patannas).

***†LEPTOMEROCORIS PUNCTATUS, sp. n.**

Long. corp. 5 millim.

Testaceous; pronotum more yellow; a band of 3 large, nearly connected black spots runs from behind each eye to the lateral angles of the pronotum, and there are 4 more conspicuous black spots arranged in a semicircle in front of the pronotum. Scutellum and corium longitudinally striped with black and testaceous: towards the extremity of the corium is a red spot on the costa bordered with yellow, and with a black spot before and behind. Membrane with the nervure black and a short black streak beyond it.

Exact locality not stated.

LIOCORIS GLABRATUS.

Liocoris glabratus, Motsch. Bull. Mosc. xxxvi. (3) p. 87, pl. ii. f. 20 (1863).

Described from Ceylon (Nura Ellia and Patannas).

***MONALOCORIS BIPUNCTIPENNIS.**

Monalocoris bipunctipennis, Walk. Cat. Hem. Het. B. M. vi. p. 159, n. 2 (1873).

Described from Ceylon.