

violaceous, the stigma and nervures black; the top of the second cubital cellule is not much more than half the length of the space bounded by the recurrent nervures. The petiole is coarsely rugosely punctured; on the sides the punctuation is coarser, more irregular, and running into reticulations.

(To be continued.)

MISCELLANEA RHYNCHOTALIA.—No. 5.

By G. W. KIRKALDY, F.E.S.

(Concluded from p. 284.)

LÆMOCORIS POSEIDON, sp. nov. ✓

Allied to *L. costæ*, Reuter, but the latter has much smaller eyes.

Clothed with sparse white pubescence. Head, pronotum, and scutellum blackish. Elytra brownish, basal third of corium (base itself excepted), a large spot on the costal margin near the apex, not reaching to the claval or membranal margin, whitish. Membrane fumate, iridescent. Antennæ brownish white, basal half of the first segment blackish brown, apical half dirty whitish. Beneath pale brownish, apex of coxæ and the rostrum whitish. Lateral margins of the sterna blackish. Eyes very large, occupying the greatest part of the head (in profile). Rostrum reaching well beyond posterior coxæ; second segment of antennæ one-fifth longer than the third, more than three times as long as the first, the latter reaching a little in front of the head. Vertex scarcely wider than an eye. Base of pronotum one-half wider than the head and eyes, twice as wide as the eyes together. Long. 3.2 mill. (with elytra), 2.5 mill. (to apex of abdomen), lat. 1 mill.

Hab. GUINEA, Addah (ex coll. Montandon). ✓

LÆMOCORIS (?) RAVENAR, sp. nov.

Clothed with fairly long, somewhat bristly, yellowish hairs. Head, pronotum, scutellum and under side, antennæ, rostrum, &c., pale flavous. Elytra pale olivaceo-flavous, a whitish band across elytra at apex of clavus, and a large whitish spot at apex of corium, not reaching to interior margins. Head (in profile) twice as long as the thickened first antennal segment, vertex a trifle wider than the eyes together. Rostrum reaching to apex of posterior coxæ. Second segment of antennæ three times as long as the first, one-sixth longer than the third. Long. 3 mill. (including elytra), lat. $\frac{3}{8}$ mill.

Hab. JAVA (ex coll. Montandon).

SHANA, gen. nov.

Belongs to *Capsaria*; distinguished from *Polymerus*, Reuter, and *Pœciloscytus*, Fieber (to which it is remotely allied) by the very distinct stink-orifices; from *Charagochilus* by the slender collar and the length of the tarsi.