

- Subg. 1.—**CAMAROCYPHUS** Reuter, 1880, Ofv. Finsk. Vet. Förh., xxii, 21. Type *nigrogularis* Reut. (= *luteus* H. S., Wanz. Ins., iii, f. 297).
- Subg. 2.—**DRYOPHILOCORIS** Reuter, 1875, Bih. Vet. Ak. Handl., iii, 30 (type fig. in Hahn. Wanz. Ins., iii, f. 235 as *flavonotatus*). Type *flavoquadrimaculatus* (de Geer), Reuter.
- Subg. 3.—**MIMOPERIDERIS** Kirkaldy n. n. = || *Perideris* Fieber, 1870, Verh. Zool. bot. Ges. Wien., xx, 249. Type *marginatus* Fieber, pl. 5, f. 6. †

Tribe 5.—**CAMPYLONEURINI**(= *Dicypharia* Reuter = *Capsaria* pt. Distant.)

- Genus 1.—**Sthenarops** Uhler, 1877, Bull. U. S. Geol. Surv., iii, 418. Type **chloris** Uhl.
- Genus 2.—**Hyaliodes** Reuter, 1875, O. V. A. F., xxxii, 83 (type fig. in Glover Ill. Ins. Hem., pl. 7, f. 18.) Type **vitripennis** (Say), Reuter.
- Genus 3.—**Engytatus** Reuter, op. c., 82 = *Neoproba* Distant, 1883, B. C. A. Het., i, 270. Type *rubescens* Distant, pl. 26, f. 6. † Type **geniculatus** Reuter.
- Genus 4.—**Campyloneura** Fieber, 1860, Eur. Hem., 67 = || *Camptoneura* Fieber, 1858, Wien. Ent. Mon., ii, 309 (type fig. in H. S. Wanz. Ins., iii, f. 268†). Type **virgula** (H. S.), Fieb. (= *pulchellus* Guér.).
- Genus 5.—**Nesidiocoris** Kirkaldy, 1902, T. E. S. London, 247. Type **volucor** Kirkaldy.
- Genus 6.—**Etorhinella** Noualhier, 1893, A. S. E. F., 16. Type **parviceps** Noualh.
- Genus 7.—**Brachyceroea** Fieber, 1858, Wien. Ent. Mon., ii, 327 = *Brachyceroea* Fieber, 1860, Eur. Hem. 77 (type fig. in Reut. Act. Soc. Fenn., xiii, pt. 3, pl. 2, f. 7 = *Dicyphus* Fieber, 1858, Wien. Ent. Mon. ii, 327 (type fig. in Reuter, op. c., pl. 3, f. 3. Type *pallidus* (H. S.)* † = † *Idolocoris* Douglas & Scott, 1865, Brit. Hem., 380. Type *pallicornis* D. & S., pl. 12, f. 5.* † Type **annulatus** (Wolff).*)
- Genus 8.—**Cyrtopeltis** Fieber, 1860, Eur. Hem., 76 (type fig. in Reuter, Act. Soc. Sci. Fenn., xiii, pt. 3, pl. 4, f. 1). Type **geniculata** Fieber.
- Genus 9.—**Macrolophus** Fieber, 1858, Wien. Ent. Mon., ii, 326 (type fig. in Reuter, op. c., pl. 4, f. 3). Type **nubilus** (H. S.).*
- Genus 10.—**Annona** Distant, 1884, B. C. A. Het., ii, 297 = || *Ania* Distant, op. c., 289. Type **bimaculata** Dist., pl. 27, f. 25. †

Tribe 6.—**CREMNORRHINI**.

- Genus 1.—**Cremonorrhinus** Reuter, 1880, Ofv. Finsk. Vet. Förh., xxii, 18 (type fig. on Reuter, Hem. Gymn. Eur., iv, pl. 1, f. 3.4† Type **basalis** Reuter.

⁴ This was probably published in the Act. Soc. Sci. Fenn., but there are no indications in the work itself that it is not a purely separate publication, and I can find no explanation in the "Zoological Record" or "Bericht der Entomologie."