

pubescence, and intermixed with more erect black pubescent hairs. Membrane and veins uniformly fumate, scarcely paler bordering apex of cuneus.

Legs: Black, tibiae and tarsi pale yellowish, basal half of hind tibiae becoming blackish, spines black but without spots at base, tips of tarsi fuscous. Venter black, brownish pubescent, sides bearing silvery scale-like pubescence.

Holotype: Female, 18 Aug., 1920, Elkhorn Creek, Carlton County, Minn. (H. H. Knight); Minn. Univ. collection.

*L. claricornis* Knight, new species.

Brownish black, antennae and tibiae pale; rostrum extending to middle of venter.

*Female*: Length 3.1 mm., width 1.47 mm. Head: Width .80 mm., vertex .36 mm.; yellowish pubescent. Rostrum, length 1.56 mm., reaching upon base of ovipositor, brownish black like the head.

Antennae: Segment i, length .19 mm., pale yellowish; ii, .77 mm., slightly thickened from base toward apex, uniformly yellowish, dusky pubescent; segments iii and iv missing.

Pronotum: Length .54 mm., width at base 1.23 mm.; brownish pubescent and intermixed on disk with closely appressed pale scale-like pubescence, propleura bearing simple pubescent hairs without any indication of scale-like pubescence. Scutellum uniformly brownish black like the pronotum, bearing scale-like pubescence intermixed with simple pubescent hairs. Sternum, pleura, and ostiolar peritreme, uniformly dark brownish, pleura bearing fine brownish pubescence only.

Hemelytra: Embolar margins slightly arcuate; uniformly brownish black, brownish pubescent and intermixed with closely appressed, pale scale-like hairs. Membrane uniformly pale brownish, scarcely paler bordering apex of cuneus, veins brownish.

Legs: Brownish black, tips of coxae and femora slightly paler; tibiae pale yellowish, spines black but without dark spot at base of each; tarsi pale, apices fuscous. Venter dark brownish, pubescence brownish.

Holotype: Female, 29 May, Lakehurst, N. J. (Wm. T. Davis); author's collection.

*L. rubidus* (Uhler).

Uhler, Hemiptera Colo., 41, 1895.

*Male*: Length 3.2 mm., width 1.5 mm.; blackish, hemelytra reddish brown with fuscous, embolium and cuneus strongly reddish; membrane uniformly fuscous; clothed with pale yellowish, closely appressed, scale-like pubescence and intermixed with more erect dusky, simple pubescent hairs; femora fusco-brownish, tinged with reddish; tibiae brownish to reddish, beset with prominent black spines. Antennae fuscous to ferruginous; segment i, length .16 mm.; ii, .64 mm., in length not equal to width of head