

Length 3 mm.; pale, smaller than *pellucida*; distinguished by the black head and fuscous basal segment of the antennae.

Food plant: Witchhazel (*Hamamelis virginiana*).

Long Island, N. Y.

Reuteria Puton.

R. irrorata (Say).

Compl. Writ., i, 346, 1859.

Length 4.3 mm., width 1.5 mm.; pale whitish, usually marked with marbled green spots; distinguished by the longitudinal black marks on antennae.

Food plants: Basswood (*Tilia*), and white oak (*Quercus*).

New York.

Noctuocoris Knight, new genus.

Arolia erect and converging at the apices as in *Orthotylus*; general aspect more nearly that of certain species of *Phytocoris*. Head distinctly oblique in female but much less so in the male; vertex ecarinate, beset with a row of erect bristles at base; an alutaceous glabrous spot each side of vertex bordering eye; clypeus large and prominent, facial angle a right angle or slightly greater; front only slightly convex. Eyes large, prominent, viewed from the side a little oblique in female, larger and more nearly vertical in male. Rostrum short, scarcely attaining hind margin of mesosternum. Pronotum moderately sloping, very little convex, sides nearly straight, forming a line which would, if projected, strike inner half of eye; callosities moderately convex, delimited behind by an impressed line which continues to an impression between the calli. Pubescence composed of two kinds of hairs, a closely appressed scale-like type, and interspersed with more erect pubescent hairs. Legs long, suggesting *Phytocoris* but femora not distinctly broader at base. Abdomen with a lateral line formed by a series of impressed glabrous spots, one on each segment (genital segments excepted) situated just below the spiracle; both sexes with abdomen reaching to about middle of cuneus. Genital claspers distinctive in type (fig. 96).

Genotype: *Orthotylus fumidus* Van Duzee.

N. fumidus (Van Duzee).

Proc. Cal. Acad. Sci., ser. 4, vi, 127, 1916.

Male: Length 5.7 mm., width 1.7 mm.; color uniformly fuscous-brown, slender apex of scutellum pale; clothed with sericeous, closely appressed, silvery and black, scale-like pubescence, and interspersed with more erect pubescent hairs, longer and more bristly at anterior margin of pronotum and on head; anterior face of femora with a more or less broken fuscous line, dorsal margin at base and frequently the posterior face with a series of dark