

Uhler, but the color characters distinguish these species at once.

♂. Length 6.2 mm., width 1.9 mm. *Head*: width 1 mm., vertex .46 mm.; vertex yellow, vertex and frons clothed with long pale sericeous pubescence. Rostrum, length 2.51 mm., reaching upon fifth ventral segment, yellowish, apex blackish. *Antennae*: segment I, length 1.57 mm., very slightly thicker on basal one-third (.143 mm. thick), beset with several pale setose hairs which in length nearly equal thickness of segment, greenish yellow; II, 2.94 mm., pale yellowish brown, paler at base, finely yellowish pubescent; III, 1.76 mm., yellowish to dusky; IV, 1.38 mm., dusky. *Pronotum*: length .96 mm., width at base 1.74 mm., width of collar .68 mm.; yellow with a tinge of greenish, median line of collar and extending between calli, and narrow basal margin of pronotum, paler. *Hemelytra*: membrane rather uniformly fumate, more brownish within areoles, veins yellowish, outside the areoles rather evenly marked with fine fuscous reticulations; clavus and corium with roseate red as described above. *Legs*: greenish yellow, apical half of hind femora somewhat darkened with fusco-brownish through which the greenish yellow color shows as small closely placed irrorate spots; tibiae pale to greenish, tips darker, spines pale brownish. Venter yellowish, pale pubescent, form of genital clasps very near *corticevivens* Knigt.

Holotype: ♂, August 19, 1924, Sierrita Mts., alt. 4,000 ft., Arizona (A. A. Nichol); author's collection. *Paratypes*: ♂, taken with type. Mr. Nichol collected this species on *Acacia*.

Phytocoris acaciae n. sp.

General color suggestive of *rufus* Van D., but distinguished by stronger pubescence combined with white and black scale-like hairs, and the conspurcate character of membrane.

♂. Length 4 mm., width 1.5 mm. *Head*: width .86 mm., vertex .30 mm. Rostrum, length 2 mm., reaching upon sixth or seventh ventral segment. *Antennae*: segment I, length .80 mm., reddish to fusco-reddish, with four or five pale spots above, spines pale; II, 1.71 mm., reddish yellow, sometimes tinged with fuscous, pale at base and with a more obscure pale band at middle; III, 1.13 mm., fuscous, paler at base; IV, .74 mm., pale fuscous. *Pronotum*: length .74 mm., width at base 1.31 mm.; disk pale fuscous, collar and basal submargin yellowish, basal margin with irregular red line and set with six tufts of black scale-like hairs. Dorsum