

6. *Psallus criocoroides* Reut. var. *Sahlbergi* Reut.

Nigro-picea, superne pilis albis squamiformibus tecta; antennis saepe articulo secundo toto flavo-ferrugineo, basi pronoti paullo breviore; hemelytris piceis vel rufescents-piceis, arcu basali cunei paullo pallidiore, membrana areola majore, apice excepto, pallida; femoribus inferne nigro-fusco-punctatis, saepe basin versus late rufo-piceis; tarsis articulo ultimo solum apice fuscescente; vertice maris oculo $\frac{2}{5} - \frac{1}{2}$ latiore. Long. $2\frac{3}{4} - 3$ mm. An species propria?

Hab. in Tunisia!, D. Prof. J. Sahlberg.

7. *Atractotomus validus* Reut.

Diagn.: Sat obscure coccineus, capite pronotoque rufo-testaceis, nonnihil fuscescens, ubique pilis brevibus aureis dense vestitus, superne subtiliter nigro-pubescent; antennis (φ) articulis duobus primis (σ) obscure coccineis, apice nigro-fuscis, tertio dimidio basali pallide flavente, dimidio apicali infuscato, secundo primo paullo crassiore, levissime fusiformi, crassitie maxima circiter quintuplo longiore; femoribus rubris, basin versus cum coxis albidis, tibiis tarsisque albido-flaventibus, his articulo ultimo fusco-nigro; hemelytris totis concoloribus, coccineis, cuneo lunula basali destituto, membrana fumata, macula ad apicem cunei posita hyalina, venis rufescents-testaceis, areola minore nigricante. Long. $\varphi 4\frac{2}{5}$ mm.

Hab. in Algeria (Bona! d. 3 maji 1896!), comm. D. E. Saunders.

Descr. Ab *A. rufo* Fieb. articulo secundo antennarum multo magis elongato, membranae macula hyalina ad apicem cunei, tibiis totis albido-flaventibus etc. mox distinguendus, ab *A. Pici* Reut. statura majore, pronoto paullo minus transverso coloreque tibiarum posticarum divergens. Caput basi pronoti circiter $\frac{3}{7}$ angustius, vertice oculo vix ma-