

*Eyes* spherical, prominent, projecting a little way beyond the anterior margin of the pronotum. *Rostrum* stout, reaching to about the second pair of coxæ.

*Thorax*: *pronotum* short, trapeziform, more than one and a half times broader on the posterior margin than long; anterior margin slightly concave; posterior margin almost straight across the scutellum, from thence to the hinder angles gently rounded. *Scutellum* triangular, equilateral, with a transverse channel near the base. *Elytra* longer than the abdomen. *Legs* moderately long and stout.

Somewhat allied to the genus *Pachylops*, Fieber; from which the flat, low crown of *Plagiotyplus*, and the shape of the face, when viewed from the side, will be found sufficient to separate it.

PLAGIOTYPLUS MACULATUS, *sp. n.* (Fieber, M. S.).

Dull green, clothed with longish semi-erect black hairs; posterior thighs sparingly punctured on the inside, next the apex. *Membrane* with numerous, small, fuscous-black spots.

*Head*: *crown* and *face* sparingly clothed with long, almost erect, stout black hairs.

*Antenna* greenish or greenish-yellow, clothed with black hairs; 1st joint green, with a few minute black punctures, in which some of the hairs are inserted; 2nd greenish-yellow, apex narrowly black; 3rd and 4th wanting. *Eyes* brown, *Rostrum* pale green, apex black.

*Thorax*: *pronotum* dull green, sparingly clothed with long, black hairs, similar to those on the head; next the anterior margin, and at a short distance from the centre, are two flat callosities, encircled by a narrow channel, and joined by a fine central one, thus having somewhat the appearance of a pair of spectacles; disc posteriorly finely crenulate. *Scutellum* slightly convex, clothed with black hairs. *Elytra* dull green. *Clavus* and *corium* somewhat thickly clothed with short, semi-depressed, black hairs. *Membrane* pale fuscous-yellow; *cells* yellowish, the entire disc and cell nerves covered with small, fuscous-black spots of irregular size, and placed at irregular intervals. *Legs* dull green or greenish-yellow; *thighs* clothed with short, black hairs. *Tibia*, on the outside, armed with a few longish, spinose, black hairs, each of which is placed in a minute, black puncture; apex brown. *Tarsi* brown. Length, 2½ lines.

Although this insect, to the best of my belief, has never been described by the late Dr. Fieber, to whom I sent it for identification, yet I retain the names he gave it, and by which it would have been known had he lived. There is only a single example, in very bad condition, taken by my friend, the Rev. T. A. Marshall, at Torla, some years ago.

Manor Park, Lee, S.E.:  
March, 1874.