quently upon the wild-rose bushes. Still, I did not find it upon the rose bushes in Clear Creek Cañon, although I swept them with much assiduity, and obtained various other insects therefrom.

HADRONEMA Uhler.

H. militaris.

Hadronema militaris Uhler, Fifth Ann. Rep. U. S. Geol. Surv. for 1871, 1872, p. 412. One specimen from Clear Creek Cañon, well up on the mountain-side, August 6.

Dr. Packard collected one specimen on July 2 at Blackhawk, Colo. It seems to be a mountain-loving species, and did not occur to me on any of the lower levels on which I used the net.

LYGUS Hahn.

1. L. lineolaris.

Capsus lineolaris Palisot-Beauv., Ins. Afr. et Amér., 187, pl. xi, fig. 7.

Almost as common in the cultivated districts of Colorado as it is in the Eastern United States. Dr. Packard found it at Golden, July 3; at Blackhawk, July 2; in the American Fork Cañon, Utah, July 22; and very common in gardens at Salt Lake City, July 25. I found it myself, in August, around Denver, near Golden, in the vicinity of Colorado Springs, and near Cañon City.

2. L. annexus.

Lygus annexus Uhler, Fifth Ann. Rep. U. S. Geol. Surv. for 1871, 1872, p. 413.

It occurred in small numbers in the mountains and cañons wherever I went. In Beaver Brook Gulch, it offered several very distinctly marked varieties, which I did not find elsewhere. Generally, the ground-color was pale olive or yellowish, marked with black or fuscous. Pale varieties were quite common near Denver on a species of *Euphorbia* with white-margined leaves.

Dr. Packard obtained it at Denver, July 28, and at Idaho, Colo., July 6.

CALOCORIS Fieber.

C. rapidus.

Capsus rapidus Say, Heteropt. New Harmony, 20, No. 4. Capsus multicolor H. Schf., Wanz. Ins., viii, 19, fig. 795.

One specimen from near Pueblo, Colo., August 10. Dr. Packard collected it in the American Fork Cañon, Utah, July 22.

RESTHENIA Amyot ET Serv.

R. confraterna.

Resthenia confraterna Uhler, Fifth Ann. Rep. U. S. Geol. Surv. for 1871, 1872, p. 411. One specimen from Beaver Brook, August 6, and another collected by Dr. Packard at Idaho, Colo., July 6.

LABOPIDEA, new genus.

Ovate, hairy, with the head broader than the pronotum, and the outer margins of the hemelytra strongly curved. Head transverse, triangular,