

tennæ; deep green, highly polished, very sparingly pubescent. Head curved posteriorly, with the neck more defined than in the preceding species, greenish-yellow, indented on the vertex, the transverse carina a little arched, with the surface in front of it a little depressed; middle of cranium with a roundish, large, black shield, which connects with the black color of the face; the whole face, under side, and base of the head polished black; tylus abruptly elevated, prominent, a little scooped out each side, not impressed at base, pubescent at tip; eyes brown, prominent. Rostrum reaching upon the intermediate coxæ, greenish; the last joint piceous; the basal joint very broad, rounded at tip; bucculæ pale. Antennæ as long as the entire hemelytra; pronotum and vertex conjoined, black, excepting the basal joint, which is green excepting the black tip and under side next the tip; the basal joint longer than the width between the eyes; second very long; third a little shorter, dusky; fourth scarcely shorter than the first, also dusky. Pronotum campanulate, longer than wide, obsolete rugose, yellow anteriorly, clothed with white, remote, pubescence, with a large, round, black spot each side, the surface distinctly sloping toward the sinuated sides; posterior margin feebly convex, the lateral angles subacutely prominent, with the margin behind them almost sinuated. Scutellum green, feebly convex, with a slightly depressed black spot near the middle at base, from which a blunt faint carina runs backward to the tip. Propleura with a large black spot reaching to the base of the coxæ; sternum dull blackish, and sometimes also the greater part of the meso- and meta-pleuræ. Coxæ green, a little dusky at base; legs green, or greenish-yellow, the tips of tibiæ and tarsi and the nails piceous. Hemelytra clear, deep green, remotely whitish-pubescent, remotely punctate, with a blackish streak on the disk of the corium; costal margin thick and prominent; membrane black or blackish, with the basal margin and the nervures pale greenish; wings blackish. Tergum black, shining; venter blackish, or with the outer margins pale, the whole surface minutely sericeous pubescent. Genital segment of the male pale piceous or yellow, the right side with a long triangular lamina extending obliquely toward the left and terminating in a long slender hook, which curves far backward.

Length to tip of venter 4-5 millimeters; to tip of membrane 5-7 millimeters. Width of base of pronotum $1\frac{1}{2}$ -2 millimeters.

Inhabits Eastern Massachusetts, New York, Pennsylvania, Maryland, in places with long grass, in June; Ohio, Southern Illinois, and Texas. Fragments of a male, a little stouter than usual, were brought from the borders of Russian America, in the far northern part of the British possessions. Missouri, C. V. Riley.

ORTHOPS Fieber.

O. scutellatus, new sp.

Form of *O. pellucidus* Fieber. Greenish-yellow when fresh. Head yellow, smooth, highly polished, impunctate; tylus black and connecting with a long black spot, which runs back to near the base of vertex;