

basal carina pale, nearly straight, narrower in the middle, and in front of this point is a faintly impressed longitudinal line; eyes dark brown, the throat and the suture between the cheeks blackish. Bucculæ black, the rostrum reaching to the posterior coxæ, greenish-yellow, with a small piceous tip. Antennæ moderately thick, yellow, tinged with piceous, the two apical joints darker, the base of third pale; second joint largest, about as long as the pronotum, the third a little longer than the basal one, and more slender than the second; fourth shortest. Pronotum with a regular curving slope toward the sides and front, smooth, polished, coarsely punctate, with a small black arc behind each callosity, a brownish cloud across the base, and a longitudinal, black, short stripe on the deflexed, punctate sides; surface a very little pubescent; posterior margin arcuated, the edge sharp, and faintly recurved; sternum blackish, the meso- and meta-pleuræ dull black, margined with yellow. Coxæ and legs greenish yellow, the knees and tips of tibiæ tinged with brown, and the apex of the tarsi with the nails piceous. Scutellum pale yellow, black at base, indistinctly and very minutely punctate, transversely rugulose, minutely pubescent. Corium dull yellow or yellowish-green, with a large brown cloud at tip and the whole clavus brownish, distinctly and closely finely punctate, minutely pubescent, the costal margin brighter yellow; cuneus dull yellow, margined behind with brown; membrane with a cloud at tip, the nervures and a tinge at base brown. Venter bright yellow, highly polished, impunctate, minutely pubescent, with a streak of brown each side, extending from the base, but not reaching to the tip.

Length to tip of venter $3\frac{1}{2}$ -4 millimeters. Width of pronotum $1\frac{1}{2}$ millimeters.

Beaten by myself from bushes in Clear Creek Cañon, August 7. It occurred singly, but was not at all rare. My short stay in that locality prevented me from collecting them in large numbers.

MERGACCELUM Fieber.

1. *M. fasciatum*, new sp.

Long-oval, yellow, stout, minutely and closely pubescent. Head broad and stout, five-angled, blunt, vertical, brownish-yellow, impunctate, broadly, obsoletely channeled transversely in front of the slender basal carina; vertex convex; tylus moderately prominent, curving inferiorly; face paler or orange, sometimes tinged with rufous; throat and bucculæ rufous. Rostrum reaching to the posterior coxæ; basal joint red, a little longer than the throat; remaining joints yellow, the second longest. Antennæ stout, reddish-yellow, paler at base; second joint a little longer than the pronotum, becoming slightly thicker toward the tip; third thicker than the preceding, and about one-third shorter; fourth thick, a little longer than the basal one. Pronotum short, minutely rugulose and obsoletely punctate, polished, pale chestnut-brown, more or less tinged with rufous; the lateral margins very oblique,