

July 9th, was given to me Mr. Henry Edwards. I have also seen specimens from Los Angeles, San Bernardino and other parts of California. The male is far more slender than the female, and has very prominent eyes and longer antennæ. The macropterous male has sometimes exceptionally long hemielytra.

*Hadronema princeps* Uhler from northern California, Oregon and Washington State, is of a narrow form, with nearly the same colors, antennæ more slender and reaching upon the cuneus, besides having a slender yellow costal border and a lunule of yellow or red on the cuneus.

*HADRONEMA MILITARIS* Uhler. One specimen from Lower California was given to me by Mr. Henry Edwards. It is also found in Arizona, Mexico, etc.

*HADRONEMA DECORATA* n. sp.

Dull black, minutely pubescent, broader than either of the other species. Head wide, moderately convex, acutely triangular in front, sometimes rufous or fulvous at base, also next the eyes, middle of front, the cheeks and throat; eyes very large and prominent, smoke-brown or black, vertex finely pubescent, minutely scabrous; antennæ long, tapering, but not abruptly narrowing, black, the second joint long, the third a little shorter, the fourth more than half the length of the third; rostrum black, sometimes fulvous at base, reaching between the middle coxæ. Pronotum about as long as wide, moderately convex, slenderly reflexed on both the anterior and lateral margins, the surface obsolete or little punctate and transversely wrinkled, the callosities transverse, small, black, sometimes enclosed by a pale band which crosses the anterior lobe, the posterior margin a little curved down, with the humeri a little tumid. Scutellum a little swollen, rufous, scabrous. Legs black, with a tinge of