

***Resthenia confraterna* Uhl.**

Colorado (Uhler, 1 and 6). Beaver Brook, August 6th (Uhler, 5). Idaho Springs, July 6th (Packard—see Uhler, 5). Colorado (Gillette—see Osborn, 1).

***Resthenia insignis* Say.**

Colorado (Uhler, 1 and 6). Colorado, as *insignis* and as *rubrivittata* (Gillette—see Osborn, 1).

North Park, July 20th (Gillette). Steamboat Springs, July 15th; Rabbit Ears Pass, July 20th (Baker).

***Resthenia insitiva* Say.**

Colorado (Gillette—see Osborn, 1).

***Resthenia maculicollis* Reut.**

Foot-hills twelve miles west of Fort Collins, June 30th (Gillette).

***Oncerometopus nigriclavus* Reut.**

Colorado (Gillette—see Osborn, 1).

Dolores, June 18th; Estes Park, July 12th (Gillette). Steamboat Springs, July 13th (Baker). Colorado Springs (Tucker).

***Oncerometopus ruber* Reut.**

Foot-hills five miles west of Fort Collins, September 1st (Gillette).

***Lopidea confluens* Say.**

Estes Park (Snow). North-east Colorado (Riley).

***Lopidea marginata* Uhl.**

Colorado (Uhler, 8).

***Lopidea media* Say.**

Foot-hills and plains, September 19th and October 4th (Carpenter—see Uhler, 6). Plentiful at Denver, Golden, Colorado Springs, and in the valley of the Arkansas near Canon City, August 5th to 18th. Occured most frequently upon the wild rose bushes (Uhler, 5). August 12th at Cottonwood Creek, Pleasant Valley, Fremont County (Cockerell, 3). West Cliff, Custer County, July 31st on *Glycyrrhiza lepidota* (Cockerell, 10). Colorado (Gillette—see Osborn, 1).

***Lopidea nigridia* Uhler n. sp.**

“Thicker and more compact than *L. media* Say, mostly dull black, pubescent, narrowly margined with rufo-fulvous or rufous on the outer border of the corium, including most of the cuneus. Head a little thicker than typical, dull black, pubescent, with a yellow spot on each side of the vertex next the eye, continued slenderly downward, a red spot on each side of the tylus, and the cheeks and sides below also red, with black sutures, the