

California Academy of Sciences (No. 328), collection of H. H. Knight, and author's collection.

34. *Orthotylus fumidus*, new species.

A large fuscous form clothed with gray pubescence. Length 6mm. to tip of membrane.

Head distinctly oblique; vertex scarcely flattened, the basal carina apparently wanting; front hardly convex, transversely striate; clypeus large, but little prominent; apex of the cheeks tumid. Eyes large, prominent; viewed from the side a little oblique, almost attaining the line of the gula. Pronotum moderately sloping, nearly flat, the sides almost rectilinear; callosities large, flat. Basal lobe of the scutellum narrowly exposed. Abdomen reaching to the tip of the cuneus. Legs long for this genus. Rostrum short, not attaining the base of the intermediate coxæ.

Color a uniform fuscous-brown, probably nearly black when fully mature; tinged with rufous on the head, apex of the pronotum, and base of the legs; sides of the venter with a black vitta. Membrane blackish, the nervures of the smaller areole pale. Whole surface clothed with short white deciduous hairs.

Described from one female example taken by me at Ft. Collins, Colo., July 28, 1898. This individual may be somewhat immature, but, while not typical of this genus, the species evidently belongs here and is included to complete the enumeration of our species, so far as they are known to me.

Type, female, in author's collection.