

Pachylops Fb.

P. punctipes (Rt.) – Jordan, several spec., 10. III. 1904, J. Sahlberg (!). – Eremian, elsewhere known from the Canary Islands, Morocco, Algeria and Tunisia.

Heterocordylus Fb.

H. tibialis (H.) – Palestine (BODENHEIMER op.cit.). – European.

Zanchius Dist.

Z. alatus Hob. – Deganya, 1 spec., 23. VII. 1958,!; Herzliya, 1 spec., 23. VII. 1958,!; Hula, 1 spec., 10. VII. 1958,!; Neve Ya'ar, 1 spec., 29. VII. 1958,!; Ramath Gan, 1 spec., 19. VII. 1958,!; Rehovot, 3 spec., 17. IX – 4. X. 1957, Swirski (!), 1 spec., 28. VII. 1958,!; Tel-Aviv, 1 spec., 24. VII. 1958,!.

On *Nymphaea* sp. in Lake Hula. Among grasses in wet biotopes. At lamps. – Syrio-Anatolian. Previously known from Turkey.

Brachynotocoris Rt.

B. viticinus Sdst. – Bat Shlomo, 7 spec., 29. VII. 1958,!; Gesher Haziv, 3 spec., 6. VIII. 1958,! Nahariya, 3 spec., 6. VIII. 1958,! – On *Vitex agnus-castus*. – Syrio-Anatolian. Previously known from Turkey and Syria.

B. cyprius E. Wgn.

Length 3.5 – 4.2 mm. Head, pronotum and scutellum yellowish white. Antennae uniformly yellowish. Eyes pale brownish grey. Elytra pale greenish; membrane pale smoky, veins pale greenish. Under surface and legs yellowish. Tip of ovipositor black.

Body gracile and flattened, $3.0 \times (\delta)$ or $2.6 \times (\varphi)$ as long as broad. Head narrower than basal width of pronotum. Vertex flattish, $1.6 \times (\delta)$ or $2.2 \times (\varphi)$ as broad as eye. Eyes relatively small. Proportions between the antennal joints $9 + 29 + 34 + 13$ (δ) or $11 + 29 + 32 + 16$ (φ) (1 unit = 0.08 mm.). Antennae with brown hairs that are longest in the 1st joint. Pronotum shorter than head, about $2.8 \times$ as broad as long, lateral margins straight; calli conspicuous, broadly separated from each other in the middle; disk flat; basal margin broadly insinuated. Elytra longer than abdomen. Hair covering of upper surface light and long. Rostrum not swollen apically. Hairs of tibiae light. Male genitalia: Right stylus (fig. 5 e) short, with two sharp apical teeth. Left stylus (fig. 5 d, f) triangular in outline, hypophysis stout with some minute teeth in the ventral margin; sensory lobe with a sharp spine.

Jerusalem, 2 spec., 13. VI. 1958,!; Palestine, 1 spec., Bodenheimer(!); Rehovot, 1 spec., 31. VII. 1957, Swirski(!); Wadi Keren near Goren, 4 spec., 6. VIII. 1958,! – On *Olea europaea*. Syrio-Anatolian. Previously recorded from Cyprus (WAGNER 1960, p. 115–116).

Platycranus Fb.

P. putoni Rt. – Beer Mashash, 1 spec., 23. VI. 1958,!; Beersheba, 1 spec., 1. VIII. 1958,!; Hadera, 2 spec., 1. VII. 1958,!; Haifa, 3 spec., 5. IV. 1904, J. Sahlberg (!), 1 spec., 12. VI. 1958,!; Herzliya, 1 spec., 4. VII. 1958,!; Ness Zionah, Petah-Tikvah, 1 spec., 1933, Carmin (!); Rehovot, 1 spec., 12. VI. 1957, Swirski (!); Revivim, 1 spec., 1. VIII. 1958,!; Wadi Rubin, 1 spec., 11. VII. 1958,!.

On *Retama raetam* on coastal dunes and in desert conditions. – Eremian, known from Morocco, Algeria, Tunisia, Egypt and Israel.