

O. laevigatus (FB.) — Tunisia: Ain Draham, 1 spec., 16—18. VIII. 1962. New to Tunisia. Holomediterranean.

Brachysteles rufescens (C.) — Tunisia: Cébola B. Ammor, 2 spec., 15. VIII. 1962. New to Tunisia. Holomediterranean.

Cardiastethus fasciiventris (G&G.) — Tunisia: Ain Draham, 1 spec., 16—18. VIII. 1962.

Microphysiidae

Myrmedobia coleoptrata (FN.) — Spanish Morocco: Gurugu, 1 spec., PARDO.

Miridae

Phytocoris minor KBM. ? — Spanish Morocco: Isaguen, 2 spec., PARDO. New to Morocco. The specimens are considerably darker than usual; the male genitalia, however, are very similar to those of European specimens. Holomediterranean.

P. algiricus RT. — Tunisia: Ain Draham, 1 spec., 16—18. VIII. 1962; Tabarka, 2 spec., 10—15. V. 1961, ECKERLEIN. On *Quercus suber*. New to Tunisia. Previously known from Algeria. Endemic.

Megacoelum quercicola n. sp.

7 mm. Pale ochraceous. Head often with 2 red median stripes continuing onto pronotum. 2nd, 3rd and 4th antennal joints usually reddish apically. Pronotum \pm darkened basally. Scutellum with a broad median dark fuscous band. Elytra ochraceous; clavus dark fuscous




Fig. 5. *Megacoelum quercicola* n. sp.: a spiculum of vesica. — *M. beckeri* (FB.): o same. — *Orthotylus junipericola* n. sp.: b—c right stylus; d left stylus. — *O. junipericola regularis* n. ssp.: e—f right stylus; g left stylus; h hypophysis of same; i chitinized appendages of vesica. — *Tinicephalus dentifer* n. sp.: j left stylus; k same, from above; l right stylus; m theca; n apex of vesica. — Orig.