

Other records: South Yemen: Aden (type locality) (Reuter 1903).

Range: East African.

Taylorilygus morosus Linnavuori

Taylorilygus morosus Linnavuori, 1975a:32.

Material: South Yemen: Jebel Jihaf, 29.IX.1937, Scott & Britton, holotype and some paratypes (BM, LC).

Range: Endemic.

Taylorilygus subdivergens Linnavuori

Taylorilygus subdivergens Linnavuori, 1974b:33–34.

Material: Yemen: Usaifira near Ta'izz, 13.XII.1937, Scott & Britton, holotype and many paratypes (BM, LC).

Range: Endemic.

**Taylorilygus pallidulus* (Blanchard)

Phytocoris pallidulus Blanchard, 1852:183.

Material: Yemen: Ghaiman near San'a slopes of Jebel Girwan, 17.II.1938, 1 ex.; San'a, 27.IX.1937, 2 exx.; Usaifira near Ta'izz, 22.XII.1937, Scott & Britton, many exx.; — South Yemen: Al Huseini near Lahej, 27.XI.1937, some exx.; Jebel Harir, 25.X.1937, Scott & Britton, 2 exx. (BM, LC).

Range: Cosmopolitan.

**Lygidolon eyrystylioides* (Wagner)

Lygus eyrystylioides Wagner, 1957:86.

Material: South Yemen: near Lahej, 13–14.VII.1963, Linnavuori, 1 ex. (LC).

Biology: On *Acacia*.

Range: Eremian.

**Orthops abessinicus* (Reuter)

Lygus abessinicus Reuter, 1903:7.

Material: Yemen: Usaifira near Ta'izz, 21.X.1937, Scott & Britton, 1 ex. (BM).

Range: Known from the Sudan, Ethiopia, Somalia, and Kenya.

Proboscidocoris punctaticollis Reuter

Proboscidocoris punctaticollis Reuter, 1905:16.

Material studied: Eritrea: Massauah, Levander, 1♂ syntype, here selected as the lectotype, and 1♂ paralectotype (Mus. Helsinki).

Often confused with the following species. Measurements: Head 0.54 × as broad as pronotum. Ocular index (♂) 1.40–1.53. 1st antennal segment 0.60–0.66 × as long as diatone, 2nd joint 3.0–3.1 × as long as 1st, 1.05–1.1 × as long as basal width of pronotum. Male genitalia as in Fig. 14a–e. Spiculum of vesica very broad, directed apicad.

Range: East African.




Fig. 14. *Proboscidocoris punctaticollis* Reuter (specimen from Massawa, Eritrea): a–b) penis; c) left style; d) right style; e) apex of right style in broad aspect.

**Proboscidocoris pilostriatus* Odhiambo

Proboscidocoris pilostriatus Odhiambo, 1959a:350–351.

Material: Yemen: Usaifira near Ta'izz, 13.XII.1937, Scott & Britton, 2 exx.; — South Yemen: Jebel Jihaf, 29.IX.1937, 3 exx.; Wadi Dareija SW of Dhala, 8.XI.1937, Scott & Britton, 2 exx. (BM, LC).

Male genitalia in Fig. 15a–c. Spiculum of vesica slender, recurved basad.

Range: East African.

Orthotylinae

**Orthotylus priesneri* Schmidt

Orthotylus priesneri Schmidt, 1939:22.

Orthotylus macrophthalmus Wagner, 1960:93–94 (Linnavuori 1986:144).

Material: South Yemen: near Lahej, 13–14.VII.1963, Linnavuori, 1 ex. (LC).

Biology: On *Acacia*.

Range: Eremian.