

**Tytthus parviceps* (Reuter)*Cyrtorhinus parviceps* Reuter, 1890:258.

Material: Yemen: Beit-el-Ghofr N of Haz, near San'a, 4.II.1938, Scott & Britton, 1 ex.; — South Yemen: Dhala, 14.IX.1937, Scott & Britton, 1 ex.; near Lahej, 9–15.VII.1963, 1 ex.; Lahej-Dhala road, 13–14.VII.1963, Linnavuori, 1 ex. (BM, LC).

Range: Intertropical.

**Yotvata picticornis* (Horvath)*Atomophora picticornis* Horvath, 1913:597.*Compsidolon acacicola* Linnavuori, 1961:18–19.

Material: South Yemen: Lahej, 9–15.VII.1963, 1 ex.; Lahej-Dhala road, 13–14.VII.1963, Linnavuori, 5 ex. (LC).

Biology: On *Acacia* and *Calligonum comosum*.

Range: Eremian.

**Plagiognathidea aervae* (Lindberg)*Malacotes aervae* Lindberg, 1958:114–115.

Material: South Yemen: near Lahej, 9–15.VII.1963, 2 exx.; Lahej-Dhala road, 13–14.VII.1963, Linnavuori, 1 ex. (LC).

Biology: On *Aerva javanica*.

Range: Eremian with a southern distribution.

Oreocapsus immundus Linnavuori*Oreocapsus immundus* Linnavuori, 1975a:100.

Material: Yemen, holotype (LC).

Range: Endemic.

**Chamaepsallus signaticornis* (Reuter)*Atomoscelis signaticornis* Reuter, 1905:20.

Material: South Yemen: al Huseini near Lahej, 29.XI.1937, Scott & Britton, 1 ex.; Lahej-Dhala road, 13–14.VII.1963, Linnavuori, many exx. (BM, LC).

Range: Eremian.

Gediocoris leptadeniae Linnavuori*Gediocoris leptadeniae* Linnavuori, 1975a:103–104.

Material: South Yemen: Anad, 9–15.VII.1963, Linnavuori, several exx. (LC).

Biology: On *Leptadenia pyrotechnica*.

Range: Eremian with a southern distribution. Known from the Sudan and Saudi Arabia.

Gediocoris vitellinus Linnavuori*Gediocoris vitellinus* Linnavuori, 1975a:104.

Material: South Yemen: Lahej-Dhala road, 9–15.VII.1963, Linnavuori, holotype and 8 paratypes (LC).

Range: Endemic.

**Campylomma plantarum* Lindberg*Campylomma plantarum* Lindberg, 1958:118–120.*Campylomma subflava* Odhiambo, 1959b:429–433.

Material: Yemen: Usaifira near Ta'izz, 13.XII.1937, Scott & Britton, 1 ex. (BM).

Range: Holosudanese. Recorded from the Cape Verde Islands, the Sudan, Uganda, and Kenya.

Campylomma hilaris Linnavuori*Campylomma hilaris* Linnavuori, 1975a:108.

Material: South Yemen: Lahej, 9–15.VII.1963, Linnavuori, 1 ex. (LC).

Biology: On *Tamarindus indicus*.

Range: Also known from the Sudan.

Campylomma acaciae Linnavuori*Campylomma acaciae* Linnavuori, 1961:22–23.

Material: South Yemen: Dhala, 14.IX.1937, Scott & Britton, 1 ex.; Lahej-Dhala road, 13–14.VII.1963, Linnavuori, several exx. (BM, LC).

Biology: On *Acacia*.

Range: Eremian.

Stenocapsus leucochilus (Reuter)*Stenarous leucochilus* Reuter, 1905:8.

Material: Yemen: near Ta'izz, on road to Mocha, 16.XII.1937, Scott & Britton, 4 exx. (BM, LC).

Range: Holosudanese.

Stenocapsus sp. near *discoidalis* (Poppius)*? Capsus discoidalis* Poppius, 1914:95.

Material: Yemen: Usaifira near Ta'izz, 22.XII.1937, Scott & Britton, 1 ex. (BM).

Auchenodes gracilis Linnavuori*Auchenodes gracilis* Linnavuori, 1975a:111.

Material: South Yemen: Wadi Kebir near Lahej, 9–15.VII.1963, Linnavuori, holotype (LC) and a paratype; Al Husaini near Lahej, 26.II.1937, Scott & Britton, several paratypes (BM, LC).

Fig. 16a–c. Closely related to *A. alboscuteolata* Puton, 1874, but longer, body much more gracile, pronotum much longer and narrower, under surface dark, scent gland orifices white, and coxae and extreme bases of femora somewhat paler (under surface in *A. alboscuteolata* largely whitish yellow and bases of femora broadly pale yellowish).

Biology: On *Tamarix*.

Range: Endemic.