

## Genus *Taeniophorus* Linnavuori

*Taeniophorus* Linnavuori 1952:36–37. Type species: *T. hyalinus* Linnavuori.

Diagnosis: Like *Camptotylidea*, but dark dotting on upper surface usually absent. Head in lateral view longer than high. Hind tarsus short and incrassate, apex of 2nd joint bluntly oblique. Pulvilli very broad, extending to near apex of claw.

Color whitish yellow, probably greenish in life. Elytra uniformly pale or with faint fuscous dots; membrane uniformly whitish hyaline. Antennae and legs pale, immaculate.

Body small, parallel-sided. Hair covering on upper surface short, pale. Head in apical view broader than high, lower part triangular in outline, head in lateral view distinctly longer than high, frons evenly sloping apicad joining without notch on to the prominent tylus, antennal pits near lower margins of eyes. Antennae in ♂ shortish and incrassate, in ♀ longish and gracile. Rostrum short, extending beyond fore coxae. Lateral margins of pronotum insinuated, calli small, distinctly elevated. Elytra longer than abdomen. Legs: tibial spines delicate, pale; hind tarsus short, about  $0.25 \times$  as long as tibia (in *Camptotylidea astarte*  $0.36 \times$ ), and incrassate, apex of 2nd joint bluntly oblique; claws gracile, pulvilli extending to near apex of claw, very broad, apex broadly rounded.

Male genitalia: Left style with coarsely dentate spine on sensory lobe. Vesica and other genitalia as in the related genera.

Distribution: Turkestan.

### *T. hyalinus* Linnavuori

Figs. 6m–s, 7v–ö, 8j–k.

*Taeniophorus hyalinus* Linnavuori 1952:37.

Type studied: Turkmenia, female allotype, J. Sahlberg, in coll. Linnavuori.

Material studied: USSR: Turkmenskaja SSR, Ahča-Kuima, 4 exx, 7.V.1974, Puchkov, in coll. Linnavuori.

Diagnosis: Characters indicated in the description of the genus.

Length 3.0–3.25 mm. Head 0.7 × as broad as basal width of pronotum. Ocular index 1.53–1.58 (♂), 2.1 (♀). Proportions between antennal joints

13:46:25:20 (♂), 11:43:25:23 (♀), 2nd joint 1.07 (♂) or 1.1 (♀) × as long as diatone, 0.75 (♂) or 0.78 (♀) × as long as basal width of pronotum.

Male genitalia in Fig. 7v–ö.

Biology: On *Eremosparton flaccidum* (Puchkov 1975:947).

Distribution: Turkestan.

Acknowledgements. I am greatly indebted to Dr. A. Jansson and Prof. H. Strümpel for allowing me to examine type material preserved in the Helsinki and Hamburg Museums, and to Prof. V. G. Puchkov, of Kiev, for providing interesting specimens of deserticolous Miridae from Central Asia.

## References

- Linnavuori, R. E. 1952: On some palearctic Hemiptera. — Ann. Entomol. Fennici 18:35–41.
- 1962: A new Atomophora species from Saudi Arabia. — Ann. Entomol. Fennici 28:69–70.
- 1971: On the genus Atomophora Rt. (Hem. Miridae). — Stuttgart. Beitr. Naturkunde 229:1–9.
- 1984: New species of Hemiptera Heteroptera from Iraq and the adjacent countries. — Acta Entomol. Fennica 44:1–59.
- 1986: Heteroptera of Saudi Arabia. — Fauna of Saudi Arabia 8:31–197.
- 1989: New species of the Miridae and Lygaeidae (Heteroptera) from the Middle East. — Ann. Entomol. Fennici 55:49–56.
- Poppius, B. 1909: Eine neue Atomophora-Art aus Sahara (Hem. Capsidae). — Ann. Entomol. Belg. 53:234.
- Puchkov, V. G. 1975: On ecology and distribution of certain little-known species of Heteroptera in the Turkmenian SSR. — Dopovidi Akad. Nauk Ukr. SSR. (B) 1975(10):946–950 (in Russian, English summary).
- 1976: New species of the genus Atomophora Reuter (Heteroptera, Miridae) inhabiting Haloxylon ssp. in the Turkmen SSR. — Dopovidi Akad. Nauk Ukr. SSR. Ser. (B) 1976(7):653–656 (in Russian, English summary).
- Reuter, O. M. 1879: Hemiptera Gymnocerata Europae. Hémiptères Gymnocerates d'Europa, du bassin de la Méditerranée et l'Asie Russe. II. — Preprint from: Acta Soc. Sci. Fenn. 1885:193–312.
- 1901: Capsidae rossicae descriptae. — Öfv. Finska Vet. Soc. Förhandl. 43:195–213.
- 1902: Capsidae novae mediterraneae IV. — Öfv. Finska Vet. Soc. Förhandl. 44:51–70.
- 1904: Capsidae novae Rossicae II. — Öfv. Finska Vet. Soc. Förhandl. 46:1–17.
- Wagner, E. 1957: Heteropteren aus Iran 1954. II. Hemiptera-Heteroptera (Fam. Miridae) (Ergebnisse der Entomologischen Reisen Willi Richter, Stuttgart im