

previously documented from the Neotropics, and this one lacks the distinctive genitalia of the Old World taxa and *Cyrtopeltocoris* Reuter from North America.

Opisthocoris
Carvalho and Costa, 1991

Opisthocoris Carvalho and Costa, 1991: 206. revised family placement

Carvalho and Costa (1991) placed *Opisthocoris*, with its single included species *O. cermelitanus* Carvalho and Costa, 1991, in the Stenodemini. Based on the habitus figure and illustrations of the pretarsus, we do not believe this taxon belongs to the Miridae, but rather to the Lygaeidae, Cyminae. Examini-

nation of specimens will be necessary to more precisely determine its identity.

Tibiopilus
Carvalho and Costa, 1993

Tibiopilus Carvalho and Costa, 1993: 207.

Carvalho and Costa (1993) placed *Tibiopilus* in the Hallopapini. Based on the description of the pronotal collar as "deprimido" (depressed) as well as the habitus figure and the illustrations of the male genitalia of the single included species *T. pedunculatus* Carvalho and Costa, we are transferring this taxon to the Phylini. No Hallopapini have been documented from the neotropics, and *Tibiopilus* appears to lack the diagnostic features of the group.

REFERENCES

- Bergroth, E.
1920. List of the Cylapinae (Hem., Miridae) with descriptions of new Philippine forms. Ann. Soc. Entomol. Belgique 60: 67–83.
- Carvalho, J. C. M.
1952. On the major classification of the Miridae (Hemiptera). (With keys to subfamilies and tribes and a catalogue of the world genera). An. Acad. Brasileira Cienc. 24: 31–110.
1955. Analecta Miridologica: miscellaneous observations in some American museums and bibliography. Rev. Chilena Entomol. 4: 221–227.
1957–60. Catalogue of the Miridae of the world, Parts 1–5. Arq. Mus. Nac. Rio de Janeiro 44: 1–158; 45: 1–216; 47: 1–161; 48: 1–384; 51: 1–194.
1984. Mirideos neotropicais, CCL: generos e especies do complexo *Pseudocarnus-Antias-Carmelus* (Hemiptera). Rev. Brasileira Biol. 44: 377–386.
1985. Mirideos Neotropicais, CCLIII: Descricoes de novos generos e especies da tribo Orthotylini Van Duzee (Hemiptera). Ibid., 45: 249–298.
1991. Mirideos neotropicais, CCCXIII: dois generos novos da tribo Phylini Douglas and Scott (Hemiptera). Bol. Mus. Paranaense Emilio Goeldi, nova serie, Zool. 6: 89–94 (1990).
- Carvalho, J. C. M., and D. L. Carpintero
1986. Mirideos neotropicais, CCLXXIII: des-
- cricoes de um novo genero & treze especies novas da Republica Argentina (Hemiptera). Rev. Brasileira Biol. 46: 607–625.
- 1991a. Miridos neotropicais CCCXLVI: descripcion de un genero y siete especies nuevas de las subfamilia Orthotylinae, con notas y correcciones taxonomicas (Hemiptera). An. Acad. Brasileira Cienc. 63: 33–42.
- 1991b. Mirideos neotropicais, CCCXLVII: novos generos e especies da Republica Argentina (Hemiptera). Ibid., 63: 201–204.
1992. Mirideos neotropicais, CCCXXXVII. descricao de doce especies novas de Orthotylinae (Hemiptera). Rev. Brasileira Biol. 51: 763–780 (1991).
- Carvalho, J. C. M., and L. A. A. Costa
1991. Mirideos neotropicais, CCCL: sobre un genero e especie curiosa de Carmo do Rio Claro, Minas Gerais (Hemiptera). An. Acad. Brasileira Cienc. 63: 205–207.
1992. Mirideos neotropicais, CCCXXXVIII: novo genero e especies de Mirinae e novas especies de Orthotylinae (Hemiptera). Rev. Brasileira Biol. 52: 199–207.
1993. Mirideos neotropicais, CCCLXXVII: dois generos e especies de Phylinae (Hemiptera). An. Acad. Brasileira Cienc. 65: 203–207.
- Cassis, G.
1986. A systematic study of the subfamily Diaphinae (Heteroptera: Miridae). Ann Arbor, Univ. Microfilms, 400 pp. [Ph.D.]