

Presentation Time: 9:30 AM-9:45 AM

SIMS OXYGEN ISOTOPE ANALYSES OF JADEITITE: TRACE ELEMENT CORRELATIONS, FLUID COMPOSITIONS, AND TEMPERATURE ESTIMATES

SORENSEN, S.S., Dept. Min. Sci, Smithsonian Inst, PO Box 37012, NMNH MRC-119, Washington, DC 20013-7012, sorena@volcano.si.edu, **HARLOW, G.E.**, Dept. Earth & Planetary Sci, AMNH, New York, NY 10024-5192, and **RUMBLE, D. III**, Geophysical Laboratory, CIW, Washington, DC 20015

Rare bodies of jadeitite (aggregates of near-end-member jadeite) represent variably deformed and recrystallized, fluid-deposited vein systems in HP/LT serpentinite-matrix mélanges. Because mineral $\delta^{18}\text{O}$ signatures can yield fluid source characteristics, we analyzed oxygen isotopes in cathodoluminescence (CL)-zoned jadeitite samples, previously analyzed for trace elements by SIMS.

Twenty $\delta^{18}\text{O}$ analyses were made for 6 jadeitites from Guatemala, California, Japan, Burma, and Kazakhstan. Separates from 2 samples and Eiler et al.'s (1997) jadeite standard were also analyzed by laser fluorination. Four samples yielded $\delta^{18}\text{O}$ from 8.18 to 9.89‰; two ranged from 4.45 to 7.07‰. Two samples are zoned in $\delta^{18}\text{O}$: in CJ-01 (Burma) bright green-CL zones (4.9‰; rich in Li, Rb, Ti, MREE, Zr) contrast with blue-CL zones (8.89 to 9.33‰, rich in Be); in sample 112552-1, (Japan), a bright green (4.45‰, rich in Li, Be, REE, Zr) zone again contrasts with red-blue zones (6.22 to 6.62‰, rich in Rb). Separates from 112552-1 yield $\delta^{18}\text{O}$ of 6.63 to 7.07‰. T-estimates based on $\delta^{18}\text{O}$ exchange between albite and jadeite range from 299° (CJ-01, Burma) to 414°C (112538, Guatemala), consistent with jadeitite-forming Ts predicted by phase equilibria and fluid inclusion data.

As has been shown for other minerals, SIMS detects large variations of $\delta^{18}\text{O}$ within (in this case, trace element- and CL-) zoned grains, which may go unnoticed in whole-rock samples or mineral separates. Globally, jadeitite-forming fluids apparently differ by as much as 5‰ in their $\delta^{18}\text{O}$ values. Jadeitite-forming fluids with such distinct $\delta^{18}\text{O}$ characteristics may have originated from different depths within subducting oceanic crust (e.g., pillow basalt versus gabbro) that underwent seafloor alteration at different temperatures, as is documented for high P/T meta-ophiolitic terrains.

[2003 Seattle Annual Meeting \(November 2–5, 2003\)](#)

Session No. 90

[Phase Relations, High P-T Terrains, P-T-ometry and Plate Pushing III: A Tribute to W.G. Ernst](#)

[Washington State Convention and Trade Center: Ballroom 6C](#)

[8:00 AM-12:00 PM, Monday, November 3, 2003](#)

© Copyright 2003 The Geological Society of America (GSA), all rights reserved. Permission is hereby granted to the author(s) of this abstract to reproduce and distribute it freely, for noncommercial purposes. Permission is hereby granted to any individual scientist to download a single copy of this electronic file and reproduce up to 20 paper copies for noncommercial purposes advancing science and education, including classroom use, providing all reproductions include the complete content shown here, including the author information. All other forms of reproduction and/or transmittal are prohibited without written permission from GSA Copyright Permissions.