

## DNA barcodes reveal species-specific mercury levels in tuna sushi that pose a health risk to consumers

Jacob H. Lowenstein, Joanna Burger, Christian W. Jeitner, George Amato, Sergios-Orestis Kolokotronis and Michael Gochfeld

*Biol. Lett.* published online 21 April 2010  
doi: 10.1098/rsbl.2010.0156

---

### Supplementary data

"Data Supplement"

<http://rsbl.royalsocietypublishing.org/content/suppl/2010/04/13/rsbl.2010.0156.DC1.html>

### References

**This article cites 16 articles, 1 of which can be accessed free**

<http://rsbl.royalsocietypublishing.org/content/early/2010/04/13/rsbl.2010.0156.full.html#ref-list-1>

### P<P

Published online 21 April 2010 in advance of the print journal.

### EXiS Open Choice

This article is free to access

### Subject collections

Articles on similar topics can be found in the following collections

[molecular biology](#) (237 articles)

[environmental science](#) (314 articles)

[health and disease and epidemiology](#) (243 articles)

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

---

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

---

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

---

# DNA barcodes reveal species-specific mercury levels in tuna sushi that pose a health risk to consumers

Jacob H. Lowenstein<sup>1,2,\*</sup>, Joanna Burger<sup>3,4</sup>,  
Christian W. Jeitner<sup>3,4</sup>, George Amato<sup>5</sup>,  
Sergios-Orestis Kolokotronis<sup>5,\*</sup>  
and Michael Gochfeld<sup>4,6,\*</sup>

<sup>1</sup>Department of Ecology, Evolution, and Environmental Biology, Columbia University, New York, NY 10027, USA

<sup>2</sup>American Museum of Natural History, Division of Vertebrate Zoology, New York, NY 10024, USA

<sup>3</sup>Division of Life Sciences, Department of Cell Biology and Neuroscience, and Consortium for Risk Evaluation with Stakeholder Participation (CRESP), Nelson Biological Laboratories, Rutgers University, Piscataway, NJ, USA

<sup>4</sup>Environmental and Occupational Health Sciences Institute, Rutgers University, Piscataway, NJ 08854, USA

<sup>5</sup>Sackler Institute for Comparative Genomics, American Museum of Natural History, New York, NY 10024, USA

<sup>6</sup>Department of Environmental and Occupational Medicine, University of Medicine and Dentistry of New Jersey, Robert Wood Johnson Medical School, Piscataway, NJ 08854, USA

\*Authors for correspondence ([jlowenstein@amnh.org](mailto:jlowenstein@amnh.org); [koloko@amnh.org](mailto:koloko@amnh.org); [gochfeld@ehsi.rutgers.edu](mailto:gochfeld@ehsi.rutgers.edu)).

**Excessive ingestion of mercury—a health hazard associated with consuming predatory fishes—damages neurological, sensory-motor and cardiovascular functioning. The mercury levels found in Bigeye Tuna (*Thunnus obesus*) and bluefin tuna species (*Thunnus maccoyii*, *Thunnus orientalis*, and *Thunnus thynnus*), exceed or approach levels permissible by Canada, the European Union, Japan, the US, and the World Health Organization. We used DNA barcodes to identify tuna sushi samples analysed for mercury and demonstrate that the ability to identify cryptic samples in the market place allows regulatory agencies to more accurately measure the risk faced by fish consumers and enact policies that better safeguard their health.**

**Keywords:** mercury; *Thunnus*; sushi; DNA barcoding; seafood labelling; epidemiology

## 1. INTRODUCTION

Accurate identification of commercial fish species has many public health and legal applications. DNA barcodes (Hebert *et al.* 2003)—short nucleotide sequences used to identify species—can serve as an important tool allowing regulatory agencies to recognize ambiguous food items that are fraudulent or hazardous (Wong & Hanner 2008; Yancy *et al.* 2008). For tuna, DNA barcodes have been used to document market substitution, and in the case of Atlantic Bluefin Tuna (*Thunnus thynnus*), meet the

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2010.0156> or via <http://rsbl.royalsocietypublishing.org>.

requirement that species protected under the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) be identifiable in trade (Lowenstein *et al.* 2009; IUCN & TRAFFIC 2010). We demonstrate one of the first applications of DNA barcoding in a human health context (Cohen *et al.* 2009) by using mitochondrial DNA to identify tuna sushi to the species level concomitant with mercury testing.

Mercury methylated by microorganisms bioaccumulates, reaching high concentrations in predatory fishes such as tuna (Morel *et al.* 1998). Excessive mercury consumption is implicated in neurodevelopmental defects including mental retardation, cerebral palsy, deafness, blindness and disarthria, and adult neuro- and cardiovascular toxicity (National Resource Council 2000). Many countries have established mercury action levels above which fish may not be sold, and have also issued advisories notifying consumers of fishes high in mercury (table 1).

Owing to relaxed international labelling requirements set by the United Nations Food and Agriculture Organization (FAO), species descriptions are often inaccurate, disputed by nations or missing (Jacquet & Pauly 2008). Many countries have ambiguous or no species-specific labelling requirements such as the US where the approved market name for all members of *Thunnus* in addition to Frigate Tuna (*Auxis thazard*), Kawakawa (*Euthynnus affinis*), Skipjack Tuna (*Katsuwonus pelamis*) and Slender Tuna (*Allothenus fallai*) is 'tuna' (FDA 2008). Tuna sushi, or *maguro* in Japanese, is made from five species sometimes specified in restaurants as bluefin tuna (*T. maccoyii*, *T. orientalis*, or *T. thynnus*), Bigeye Tuna/ahi (*T. obesus*) or Yellowfin Tuna/ahi (*T. albacares*; Lowenstein *et al.* 2009). Because of overlap in appearance and taste (Catarci 2005), molecular identification is one of the most precise methods for identifying tuna in the marketplace (Lowenstein *et al.* 2009; Viñas & Tudela 2009). In the context of mercury analysis, DNA barcodes enabled us to determine which species warrant inclusion in consumer advisories or trade restrictions, and whether the data used by health agencies reflect accurately the mercury threat faced by consumers.

## 2. MATERIAL AND METHODS

We tested the mercury content of 100 tuna sushi samples from 54 restaurants and 15 supermarkets collected from October 2007 to December 2009 in New York, New Jersey, and Colorado. The New York Times collected 20 samples, and we collected the rest. We identified them using nucleotide characters and BLASTN against NCBI GenBank (see the electronic supplementary material, table S1) using the cytochrome *c* oxidase subunit I (*cox1*) gene sequence, following the methodology detailed in Lowenstein *et al.* (2009). Because the three closely related species of bluefin are often not differentiated in global trade (Catarci 2005) we pooled these data into one category for the mercury analysis. We further categorized samples according to whether they were sold as lean red tuna (*akami* in Japanese) or fatty tuna (*toro*) because mercury and lipid concentrations are inversely proportional in tuna (Balshaw *et al.* 2008a).

To measure total mercury, a 2 g (wet weight) subsample of fish tissue was digested in trace metal grade nitric acid (Fisher Chemical) in a microwave (CEM, MDX 2000), using a digestion protocol of three stages of 10 min each under 50, 100 and 150 pounds per square inch (3.5, 7 and 10.6 kg cm<sup>-2</sup>) at 80 per cent power. Digested samples were subsequently diluted to 25 ml with deionized water. All laboratory equipment and containers were washed in 10 per cent HNO<sub>3</sub> solution and rinsed with deionized water prior to each use.

Table 1. Mercury advisory levels set by regulatory agencies. (The weekly maximum level recommended by the Food and Agriculture Organization of the United Nations and the World Health Organization for women of childbearing age is equivalent to  $0.2 \mu\text{g kg}^{-1}$  body weight per day.)

agency	mercury advisory level	reference
European Commission	1.0 ppm	European Commission (2008)
Food and Agriculture Organization	$1.6 \mu\text{g kg}^{-1}$ body weight per week	Codex Alimentarius Commission (1995)
Health Canada	1.0 ppm	Health Canada (2007)
Japanese Ministry of Health	0.4 ppm	Yamashita (2005)
US Environmental Protection Agency	$0.1 \mu\text{g kg}^{-1}$ body weight per day	EPA (1997)
US Food and Drug Administration	1.0 ppm	FDA (2000)
World Health Organization	$1.6 \mu\text{g kg}^{-1}$ body weight per week	Codex Alimentarius Commission (1995)

Table 2. Total mercury (Hg) content in tuna sushi samples. (Data from samples identified as one of the three species of bluefin (*T. maccoyii*,  $n = 7$ ; *T. orientalis*,  $n = 4$ ; *T. thynnus*,  $n = 18$ ), were pooled into a single category. *Akami* is the Japanese word for lean red tuna, and *toro* for fatty tuna. Total mercury (ppm) varied significantly across sample categories (one-way ANOVA:  $F_{4,95} = 11.81$ ,  $p < 0.0001$ ). Categories assigned as 'a' were significantly different from those assigned 'b' (Tukey's multiple comparison test). The mean dose was calculated for the default weight of a 60 kg adult woman (WHO 1972; Health Canada 2007) consuming a single order.)

sample category	total Hg (ppm)						mean dose of total Hg ( $\mu\text{g kg}^{-1}$ body weight per day)	sample mass (g; mean $\pm$ s.e.)	sample size
	mean	median	s.d.	min	max	assignment			
Bigeye Tuna akami	0.871	0.794	0.393	0.336	1.716	a	0.344	$22.48 \pm 2.843$	36
Bigeye Tuna toro	0.989	0.685	0.716	0.365	2.254	a	0.351	$20.82 \pm 2.941$	9
bluefin tuna akami	1.043	1.028	0.478	0.368	1.916	a	0.180	$12.09 \pm 2.046$	10
bluefin tuna toro	0.385	0.307	0.244	0.166	1.027	b	0.123	$21.18 \pm 2.428$	19
Yellowfin Tuna akami	0.474	0.435	0.294	0.095	1.377	b	0.164	$18.34 \pm 2.823$	26

Mercury was analysed by cold vapour technique using a Perkin Elmer FIMS-100 mercury analyser, with an instrument detection level of  $0.004 \mu\text{g g}^{-1}$  and a method detection level of  $0.010 \mu\text{g g}^{-1}$ . All samples were tested twice, and 15 samples with the highest mercury levels were tested three times. For 98 of the samples results from all runs were within 5 per cent, and two samples within 10 per cent. As a control, we used the National Institute of Standards and Technology dogfish muscle trace metal reference material (DORM-2) alongside the samples and our results were always within the total mercury certificate range (4.38–4.90 ppm). For sushi-grade tuna, approximately 97 per cent of total mercury is methylmercury (Hight & Cheng 2006). The mercury analysis was carried out at Rutgers University and the genetic identification at the American Museum of Natural History with no prior knowledge of sample identity or mercury concentration.

Statistical analyses were performed on GRAPHPAD PRISM v. 5.00b for Mac OSX ([www.graphpad.com](http://www.graphpad.com)). Before using parametric tests, we performed a  $\log(x + 1)$  transformation on all mercury data used in this study and assessed conformation to a normal distribution using the D'Agostino–Pearson omnibus test ( $\alpha = 0.05$ ). We assessed homogeneity of variances for the one-way ANOVA using Bartlett's test, as well as for comparisons of two groups using an *F*-test ( $\alpha = 0.05$ ).

### 3. RESULTS

Mercury concentrations varied significantly across sample categories (one-way ANOVA:  $F_{4,95} = 11.81$ ,  $p < 0.0001$ ; table 2). The mercury levels in bluefin *akami* and all Bigeye Tuna samples were significantly higher compared with bluefin *toro* and Yellowfin Tuna *akami*. The mean mercury concentrations of all samples exceed the concentration permitted by Japan (Yamashita *et al.* 2005), and the maximum daily

consumption considered safe by the US Environmental Protection Agency (EPA 1997). Mean mercury levels for bluefin *akami* exceed those permitted by the US Food and Drug Administration (2000), Health Canada (2007) and the European Commission (2008). On average, one order of Bigeye Tuna sushi—the species used most often for sushi (Catarci 2005)—exceeds the safe maximum daily dose recommended by Health Canada (2007) and the safe limit established by the World Health Organization and FAO for women of childbearing age (Codex Alimentarius Commission 1995).

As documented previously for Southern Bluefin Tuna (Balshaw *et al.* 2008a), we found significantly less mercury in bluefin *toro* than in *akami* ( $t = 5.109$ ,  $p < 0.0001$ ), but no significant difference for Bigeye Tuna ( $t = 0.363$ ,  $p = 0.717$ ). We found no significant difference in bluefin mercury levels comparing data from a study (Storelli *et al.* 2002) with greater sampling ( $n = 161$ , mean = 1.18 ppm, s.d. = 0.85) to our bluefin *akami* results (Mann–Whitney *U*-test,  $p = 0.59$ ). The total mercury levels we found in Yellowfin Tuna sushi was significantly higher (Mann–Whitney *U*-test,  $p = 0.0236$ ) than in samples obtained by the FDA (2004), as was the case for Bigeye Tuna (*t*-test with Welch's correction,  $t = 2.549$ ,  $p = 0.0162$ ; figure 1). Finally, we found that the concentration of total mercury was also higher in our samples sold in restaurants compared with supermarkets ( $t = 3.249$ ,  $p = 0.0018$ ; figure 1).




Figure 1. (a) Total mercury concentration (ppm; mean  $\pm$  s.e.) in Bigeye Tuna samples (squares) and Yellowfin Tuna (circles) collected by the US Food and Drug Administration (FDA; unfilled) and sushi samples collected for this study (filled). The FDA lacks data on mercury levels in bluefin tuna. Sushi samples represent both *akami* (lean red tuna in Japanese) and *toro* (fatty tuna in Japanese). (b) Mean total mercury in *akami* sushi samples sold in supermarkets (unfilled diamonds) and restaurants (filled diamonds) for all species. No *toro* was found in supermarkets.

#### 4. DISCUSSION AND CONCLUSIONS

Our results demonstrate the use of DNA barcodes to enable regulatory agencies to identify unknown and potentially hazardous samples. A multi-locus genetic species identification method was recently proposed for tuna (Viñas & Tudela 2009), and while we agree that multi-locus approaches perform better in cases of introgressive hybridization, this discussion does not have a negative impact on our findings presented here.

Mercury concentrations in tuna are positively correlated with body size (Storelli *et al.* 2002; Yamashita *et al.* 2005), and larger individuals are more likely to be sushi-grade and valued the highest (Catarci 2005). The finding that the mercury levels in Bigeye Tuna *akami* and *toro* were not significantly different may be owing to the fact that premium Bigeye *toro* cuts on average have half the fat content of bluefin (Shimamoto *et al.* 2003; Balshaw *et al.* 2008a) and because larger fish typically have more belly fat and are preferentially selected for *toro*. Furthermore, whereas thousands of tons of bluefin per year are fattened in farms prior to export (Catarci 2005), which can also reduce mercury (Balshaw *et al.* 2008b), the vast majority of Bigeye Tuna are harvested directly from the wild. Because the mercury concentrations found in our sushi were significantly higher than levels documented by the Food and Drug Administration (FDA) (figure 1), this could reflect that our samples came from larger fish (the FDA lacks bluefin data). We found significantly lower mercury levels in supermarket sushi (figure 1) because samples were dominated (77%) by Yellowfin Tuna, which comprised a minority of restaurant samples (22%;  $\chi^2 = 18.14$ ,  $p < 10^{-4}$ ) and was found to be the species with the lowest mercury concentration (table 2). By allowing for the direct measurement of samples collected from the marketplace, DNA barcoding has the potential of revealing mercury measurements more reflective of the threat faced by consumers

allowing for the enactment of policy that better safeguards consumer health.

Our results suggest health agencies should consider adding Bigeye and bluefin tuna to mercury advisories. For instance, the mercury levels in these species are within the bounds of fish the FDA and EPA advise pregnant or nursing women and children to avoid entirely (EPA & FDA 2004), and thus these tunas should be included in the advisory. Consumers could make more informed health decisions if the FDA, and regulatory agencies in other nations, enforced market-specific names for species high in mercury.

We thank the New York Times for samples, and the Alfred P. Sloan Foundation, the Richard Lounsbery Foundation, and Columbia University for funds. Mercury analyses were partially funded by NIEHS Center Grant P30ES005022. We are grateful for the comments and constructive criticisms from two anonymous reviewers that helped to improve the manuscript.

- Balshaw, S., Edwards, J., Ross, K. & Daughtry, B. 2008a Mercury distribution in the muscular tissue of farmed southern bluefin tuna (*Thunnus maccoyii*) is inversely related to the lipid content of tissues. *Food Chem.* **111**, 616–621. (doi:10.1016/j.foodchem.2008.04.041)
- Balshaw, S., Edwards, J. W., Ross, K. E., Ellis, D., Padula, D. J. & Daughtry, B. J. 2008b Empirical models to identify mechanisms driving reductions in tissue mercury concentration during culture of farmed southern bluefin tuna *Thunnus maccoyii*. *Mar. Pol. Bull.* **56**, 2009–2017. (doi:10.1016/j.marpolbul.2008.08.007)
- Catarci, C. 2005 The world tuna industry: an analysis of imports and prices, and of the combined impact on catches and tuna capacity. In *Management of tuna fishing capacity: conservation and socio-economics*, vol. 2 (eds W. H. Bayliff, J. I. L. Moreno & J. Majkowski), pp. 279–321. Madrid, Spain: FAO.
- Codex Alimentarius Commission 1995 Codex general standard for contaminants and toxins in food and feed

- (CODEX STAN 193–1995). Joint FAO/WHO food standards programme.
- Cohen, N. *et al.* 2009 Public health response to puffer fish (Tetrodotoxin) poisoning from mislabeled product. *J. Food Prot.* **72**, 810–817.
- EPA 1997 Mercury study report to congress. Washington, DC: U.S. EPA Office of Air Quality Planning and Standards and Office of Research and Development.
- EPA & FDA 2004 What you need to know about mercury in fish and shellfish: advice for women who might become pregnant, women who are pregnant, nursing mothers, young children. EPA-823-R-04-005. See <http://www.fda.gov/Food/FoodSafety/Product-SpecificInformation/Seafood/FoodbornePathogensContaminants/Methylmercury/ucm115662.htm>.
- European Commission 2008 Commission regulation (EC) No. 629/2008 of 2 July 2008 amending Regulation (EC) No 1881/2006 setting maximum levels for certain contaminants in foodstuffs. *Off. J. EU* **51**, 4–10.
- FDA 2000 Guidance for industry: action levels for poisonous or deleterious substances in human food and animal feed. See <http://www.fda.gov/Food/Guidance-ComplianceRegulatoryInformation/GuidanceDocuments/ChemicalContaminantsandPesticides/ucm077969.htm>.
- FDA 2004 Mercury concentrations in fish: FDA monitoring program (1990–2004). See <http://www.fda.gov/Food/FoodSafety/Product-SpecificInformation/Seafood/FoodbornePathogensContaminants/Methylmercury/ucm191007.htm>.
- FDA 2008 Seafood complete list. U.S. Food and Drug Administration. See [http://www.accessdata.fda.gov/scripts/SEARCH\\_SEAFOOD/index.cfm?other=complete](http://www.accessdata.fda.gov/scripts/SEARCH_SEAFOOD/index.cfm?other=complete).
- Health Canada 2007 Human health risk assessment of mercury in fish and health benefits of fish consumption. See [http://www.hc-sc.gc.ca/fn-an/pubs/mercure/merc\\_fish\\_poisson-eng.php](http://www.hc-sc.gc.ca/fn-an/pubs/mercure/merc_fish_poisson-eng.php).
- Hebert, P. D. N., Cywinska, A., Ball, S. L. & DeWaard, J. R. 2003 Biological identifications through DNA barcodes. *Proc. R. Soc. Lond. B* **270**, 313–321. (doi:10.1098/rspb.2002.2218)
- Hight, S. C. & Cheng, J. 2006 Determination of methylmercury and estimation of total mercury in seafood using high performance liquid chromatography (HPLC) and inductively coupled plasma-mass spectrometry (ICP-MS): Method development and validation. *Anal. Chem. Acta* **567**, 160–172. (doi:10.1016/j.aca.2006.03.04)
- IUCN & TRAFFIC 2010 In *IUCN/TRAFFIC analysis of the proposals to amend the CITES appendices*. IUCN, the International Union for Conservation of Nature. Prepared by IUCN Species Programme, SSC and TRAFFIC for the Fifteenth Meeting of the Conference of the Parties to CITES. Gland, Switzerland.
- Jacquet, J. & Pauly, D. 2008 Trade secrets: renaming and mislabeling of seafood. *Mar. Pol.* **32**, 309–318. (doi:10.1016/j.marpol.2007.06.007)
- Lowenstein, J., Amato, G. & Kolokotronis, S. 2009 The real *maccoyii*: identifying tuna sushi with DNA barcodes: contrasting characteristic attributes and genetic distances. *PLoS ONE* **4**, e7866. (doi:10.1371/journal.pone.0007866)
- Morel, F. M. M., Kraepiel, A. M. L. & Amyot, M. 1998 The chemical cycle and bioaccumulation of mercury. *Annu. Rev. Ecol. Syst.* **29**, 543–566. (doi:10.1146/annurev.ecolsys.29.1.543)
- National Resource Council 2000 *Toxicological effects of methylmercury*. Washington, DC: National Academy Press.
- Shimamoto, J., Hasegawa, K., Hattori, S., Hattori, Y. & Mizuno, T. 2003 Non-destructive determination of the fat content in glazed bigeye tuna by portable near infrared spectrophotometer. *Fish. Sci.* **69**, 1247–1256. (doi:10.1111/j.0919-9268.2003.00752.x)
- Storelli, M. M., Stuffer, R. G. & Marcotrigiano, G. O. 2002 Total and methylmercury residues in tuna-fish from the Mediterranean sea. *Food Add. Cont.* **19**, 715–720. (doi:10.1080/02652030210153569)
- Viñas, J. & Tudela, S. 2009 A validated methodology for genetic identification of tuna species (genus *Thunnus*). *PLoS ONE* **4**, e7606. (doi:10.1371/journal.pone.0007606)
- WHO 1972 *World health organization food additive series*. vol. 52. *Methylmercury (addendum)*. Geneva, Switzerland: WHO.
- Wong, E. & Hanner, R. 2008 DNA barcoding detects market substitution in North American seafood. *Food Res. Int.* **41**, 828–837. (doi:10.1016/j.foodres.2008.07.005)
- Yamashita, Y., Omura, Y. & Okazaki, E. 2005 Total mercury and methylmercury levels in commercially important fishes in Japan. *Fish. Sci.* **71**, 1029–1035. (doi:10.1111/j.1444-2906.2005.01060.x)
- Yancy, H. *et al.* 2008 Potential use of DNA barcodes in regulatory science: applications of the regulatory fish encyclopedia. *J. Food Prot.* **71**, 210–217.