

- 1917 ————— Van Duzee, Univ. Cal. Pub. Ent. 2:390 (as type)
 1923 ————— Butler, Biol. Brit. Hem.: 486 (biology)
 1933 ————— Stichel, Illus. Best. Deut. Wanz. 8:229 (key)
 1935 ————— Hedicke, Tierw. Mittel. Het. 4(3):57 (key)
 1939 ————— f. *striola* Wagner, Jahrb. Nass. Ver. Naturkd. 86:45 (note)
 1943 ————— China, Gen. Names Brit. Ins. 8, Hem.: 265 (as type)
 1951 ————— Kiritshenko, Het. Eur. USSR.: 177 (key)
 1952 ————— Wagner, Tierw. Deut. 41, Blindw.: 122 (key, descr.)
 1952 ————— Carvalho, An. Acad. Brasil. Ci. 24(1):80 (as type)
 1953 ————— Southwood, Trans. R. Ent. Soc. London 104(11):431, figs.
 (key)
- necopinus* Van Duzee N. America
 1916 *Orthotylus necopinus* Van Duzee, Proc. Cal. Acad. Sci. (4)6:125 (n. sp.) Ill.
 1917 ————— (*Diommatus*) *necopinus* Van Duzee, Univ. Sal. Pub. Ent. 2:395 Minn.
 (cat.) N.H.
 1923 ————— *necopinus* Knight, Conn. Nat. Hist. Surv. Bul. 34:512 (key), 517 N.Y.
 (descr.), fig. 87 Ont.
 1926 ————— Blatchley, Het. E.N. Amer.: 861 (descr.)
 1941 ————— Knight, Ill. Nat. Hist. Surv. Bul. 22(1):98, 99, 105
 (key, descr.) fig. 128
- neglectus* Knight N. America
 1923 *Orthotylus neglectus* Knight, Conn. Nat. Surv. Bul. 34:515 (n. sp.), 513 (key), Conn.
 fig. 84 Ill.
 1926 ————— Blatchley, Het. E.N. Amer.: 865 (descr.) Ind.
 1941 ————— Knight, Ill. Nat. Hist. Surv. Bul. 22(1):98, 99, 105 (key, N.S.
 descr.) fig. 128 N.Y.
- nigrinasi* Van Duzee see *Ilnacora* Reuter
- nigripes* Reuter see *Labopidea* Uhler
- nigropilosus* Lindberg see *Melanotrichus* Reuter
- notabilis* Knight N. America
 1927 *Orthotylus notabilis* Knight, Can. Ent. 59:176 (n. sp.) Ia. Kans.
 1941 ————— Knight, Ill. Nat. Hist. Surv. Bul. 22(1):97, 98, 100 (key, Minn.
 distr.) Sask.
 1949 ————— Froeschner, Amer. Mid. Nat. 42(1):142, 171 (key, note) S.E.
- nyctalis* Knight N. America
 1927 *Orthotylus nyctalis* Knight, Can. Ent. 59:181 (n. sp.) Ia. Ill. Minn.
 1941 ————— Knight, Ill. Nat. Hist. Surv. Bul. 22(1):98, 99, 104 N.Y.
 (key, distr.) fig. 128 Wiss.
- obscurus* Reuter Austria
 * 1835 *Phytocoris mutabilis* Hahn nec Fallen, Wanz. Ins. 2:137, fig. 223 (descr.) (syn. France
 by Reuter, Hem. Gymn. Eur. 3:344, 1883) Germany
 1861 *Brachyarthrum pinetullum* Fieber nec Fallen, Eur. Hem.: 301 Sweden
 1875 *Orthotylus obscurus* Reuter, Pet. Nouv. Ent. 1:(138):551 (n. sp.) Yugoslavia
 * 1875 ————— *fuscescens* Reuter nec Kirschbaum, Bih. K. Sv. Vet. Akad. Handl. Rumania
 3(1):33 (syn. by Reuter, Hem. Gymn. Eur. 3:344, 1883)
 1883 ————— *obscurus* Reuter, Hem. Gymn. Eur. 3:344, 549 (descr., key)
 1890 ————— Atkinson, Jour. Asiatic Soc. Bengal 58(2):138 (cat.)
 1908 ————— Hueber, Syn. Blindw. 2:156, 162 (key, descr.)
 1909 ————— Reuter, Acta Soc. Sci. Fenn. 36(1):76 (host)
 1910 ————— Oshanin, Verz. Pal. Hem.: 837 (cat.)
 1933 ————— Stichel, Illus. Best. Deut. Wanz. 8:227 (key)
 1935 ————— Hedicke, Tierw. Mittel. Het. 4(3):56 (key)
- * *obsoletus* Fieber see *adenocarpi* Perris
- ochrotrichus* Fieber G. Britain
 1865 *Litosoma ochrotrichus* Douglas & Scott, Brit. Hem.: 342 (descr.) Morocco
 1864 *Orthotylus ochrotrichus* Fieber, Wien Ent. Monat. 8:330 (n. sp.) Spain
 1873 *Leptomerocoris ochrotrichus* Walker, Cat. Het. 6:133 (list)
 * 1883 *Orthotylus propinquus* Reuter, Hem. Gymn. Eur. 3:356, 551, pl. 5, fig. 17
 (descr., key) (syn. by Oshanin, Verz. Pal. Hem.: 841, 1910)