

Rostrum reaching middle coxae.

Genitalia: Aedeagus (fig. 8a) with vesica showing typical secondary gonopore. Left clasper (fig. 8b) as shown in figure. Right clasper (fig. 8c) small, as illustrated.

FEMALE. Similar to male in color. Length 2.8 mm., width 1.2 mm.




FIGURE 8. *Campylomma citrina* Carvalho, new species: a, Aedeagus; b, Left clasper; c, Right clasper.

HOLOTYPE. Male, Galápagos, Archipelago, Bella Vista, Santa Cruz Island, II-4-64 (Usinger).

ALLOTYPE. Female, same data as holotype.

PARATYPES. Three males, same data as type.

ADDITIONAL SPECIMENS. SANTA CRUZ: 6 ♂♂, 18 ♀♀, Bella Vista, II-4-1964 (Usinger). FLOREANA: 1 ♂, 1 ♀, Wittmer Farm, II-15-1964 (Usinger); 1 ♀, same data, II-18-1964.

This species approaches *Campylomma verbasci* (Meyer) and *Campylomma unicolor* Poppius in the color of antennae but differs from them by the general color of the body and by the structure of the aedeagus. The spots of posterior femora are also characteristic of this species.

Genus PSALLUS Fieber
Key to Galápagos Species

1. Rostrum reaching beyond apex of hind coxae, attaining genital segment (male) or 7th to 8th segment (female) *P. longirostris*