

B: 1 4.8 w 1.5 H: 1 0.3 v 0.33 A: I 0.7
II 2.0 III-IV mutilated P: 1 0.6 w 1.3

278. *villosa* CALOCORISCA 1884: 280, Pl 26: 17 —
CALOCORISCA *villosa* L — ♀, V. de Chiriqui,
8000 ft. Champion. P — 4. (Altitude of V.
de Chiriqui locality cited in description only
as 4000-6000 ft. two paralectotypes have
such data, but perhaps the altitude difference
was overlooked by Distant, so we
accept the specimen labelled "Type" as
lectotype.

B: 1 9.1 w 4.0 H: 1 0.6 w 1.5 v 0.46 A:
I 0.7 II 2.5 III 0.9 IV mutilated P: 1 1.6
w 3.1

279. *villosus* NEOFURIUS 1884: 292, Pl 28: 13 —
KNIGHTOCORIS *villosus* (Distant): L — ♀, V.
de Chiriqui, 2-3000 ft. Champion. P — 0.

B: 1 4.8 w 2.1 H: 1 0.7 w 0.9 v 0.54 A:
I 0.6 II 0.8 III 0.6 IV 0.6 P: 1 1.1 w 1.7
C: 1 0.85 w 0.62

280. *villosa* DEMARATA 1884: 303 Pl 29: 14 — CERA-
TOCAPSUS *villosus* (Distant): L — ♀, Gua-
temala City, Champion. P — 9.

B: 1 3.1 w 1.5 H: 1 0.3 w 0.7 v 0.36 A:
I 0.2 II 0.6 III 0.4 IV 0.3 P: 1 0.6 w 1.2
C: 1 0.36 w 0.58

281. *vinaceus* LYGUS 1884: 273-274, Pl 26: 11 —
PROBA *vinacea* (Distant): L — ♂, Toton-
capam, 85-10500 ft. Champion. P — 1.

B: 1 7.3 w 3.0 H: 1 0.8 w 1.3 v 0.39 A:
I 0.7 II 2.1 III 1.1 IV 0.7 P: 1 1.2 w 2.4
C: 1 0.71 w 0.62

282. *vitreus* NEOCARNUS 1884: 289, Pl 29: 1 — HYA-
LIODES *vitreus* (Distant): L — sex unknown,
Bugaba, Panama. Champion. P — 1.

B: 1 4.4 w 1.4 H: 1 0.4 w 0.7 v 0.32 A:
I 0.7 II 1.2 III 0.4 IV 0.5 P: 1 0.8 w 1.1
C: 1 0.66 w 0.36

283. *vulgris* ORNANDES 1893: 448-449, Pl 39: 13 —
CHLAMYDATUS *associatus* (Uhler): L — ♀,
Teapa, Tabasco. March. H. H. Smith.
P — 1.

B: 1 3.0 w 1.4 H: 1 0.4 w 0.7 v 0.42 A:
I 0.2 II 0.7 III 0.3 IV 0.3 P: 1 0.5 w 1.1
C: 1 0.40 w 0.49

284. *vultuosus* ECCRITOTARSUS 1893: 440, Pl 38: 13
— PACHYPODA *vultuosa* L — ♂, Frontera,
Tabasco. Jan. H. H. Smith. P — 1.

B: 1 8.7 w — H: 1 0.4 w 0.8 v 0.50 A:
I 0.4 II 0.8 III 0.5 IV 0.5 P: 1 0.6 w 1.2
C: 1 0.66 w 0.38

SUMMARY

All of the 284 species of Miridae originally described by W. L. Distant in the "Biologia Centrali Americana" (1883, 1884 and 1893) are listed in alphabetical order together with the present binomen by which each of his nominal species is currently known. Type material was traced for 277 of these species and its type status assessed. For all species except one of these it was possible either to recognize a holotype or to designate a lectotype. Measurements of 14 parameters and the sex of all primary types are given, where these could be determined. All type material of Distant's species still known to exist is listed. It is preserved in the British Museum (Natural History), London, the Zoologisches Museum of the Humboldt-Universität, Berlin and the Naturhistoriska Riksmuseet, Stockholm.

REFERENCES

- CARVALHO, J. C. M., 1952, Neotropical Miridae, L: On the present Generic Assignment of the Species in the Biologia Centrali Americana (Hemiptera). *Bol. Mus. Nac. R. Jan.*, n. s., Zoologia, 118: 17 p.
- DISTANT, W. L., 1880-1893, *Biologia Centrali Americana*. Insecta. Rhymchota. Hemiptera-Heteroptera, vol. 1: 1-302 (1880-1884); Supl.: 304-462 (1884-1893), 39 pls., London.