

***Pseudoloxops vitiensis* (Kirkaldy, 1908)**

Carvalho, 1955

(Figs. 7-11)

Tichorhinus vitiensis Kirkaldy, *Sydney Proc. Linn. Soc.*, 33: 377.*Pseudoloxops vitiensis* Carvalho, *Rev. Chil. Ent.*, 4: 224, 1955; Id. Carvalho, *Arq. Mus. Nac. R. Jan.*, 47: 128, 1958.


Characterized by its color and structure of male genitalia.

Male: Length 3.6 mm, width 1.2 mm.

Head: Length 0.2 mm, width 0.7 mm, ver-

tex 0.36 mm. *Antenna*: Segment I, length 0.4 mm; II, 1.6 mm; III, 0.6 mm; IV, 0.6 mm. *Pronotum*: Length 0.4 mm, width at base 1.0 mm. *Cuneus*: Length 0.44 mm, width at base 0.28 mm.

General coloration reddish with pale yellow areas; segment I and base of segment II of antennae, head, pronotum (except semilunar roundish central pale yellow spot on disc), scutellum (except pale basal portion), clavus (except large median pale spot), embolium and corium (except apical



Pseudoloxops vitiensis (Kirkaldy) — Fig. 7: Male, holotype; fig. 8: penis; fig. 9: left paramere; figs. 10 and 11: right paramere.