

Cyrtocapsidea irrorata Reuter, 1907

Four females: 2 ex. Bilimek, Orizaba, 5 Mai; Bilimek, Mexico, 1871. Segment I of antenna black (pale at extreme apex); segment II pale, black on apical third; segments III and IV pale at base; scutellum tumid, pale apically; cuneus black, apex pale; disc of pronotum with three dark longitudinal fasciae. Small size. Belongs to *Derophthalma* Berg.

Cyrtocapsus rostratus Reuter, 1908

One specimen, female, Tombillo, Otto, 1895. Pubescence woolly, silvery, hind margin of pronotum brown, apex of corium with a pale spot, legs and antennae pale, vertex and frons with a longitudinal pale fascia, hairs whitish; head pointed in front, segment II of antenna long and slender.

Dichroocoris coerulipennis Reuter, 1908

One specimen, male, Mexico, Bilimek, Orizaba, 5 Mai. Color bluish black brilliant metallic with head, pronotum, mesoscutum and base of hemelytra rufescent (reddish lutescent), coxae and legs pale.

Eccritotarsus delicatus (Reuter, 1907) Hsiao, 1947

One specimen, Stieglmayr, Rio Grande do Sul. Color pale yellow to whitish, pronotum tending to lutescent, antennae, eyes, apex of scutellum, apex of clavus, transversal fascia on corium and apex of hind femur black; apical portion of embolium, extreme apex of corium, apex of cuneus internally, nervure and small spot at base of areola fuscous to black.

Eccritotarsus discipennis Stål, 1860

Blumenau, Hetschko, 1889, 1 specimen. Head pale, pronotum and scutellum black; hemelytra pale, claval commissure, apex of clavus, transverse fascia on corium at level of apex of clavus, cuneus and membrane pale; legs and antenna pale.

Eccritotarsus nigrocruciatus pallidicornis
Reuter, 1907

Rio Grande do Sul, Stieglmayr, 1 specimen. Black, basal half of corium and embolium and two basal thirds of cuneus pale, nervures of membrane and segments I and II of antenna brown, apex of segment II black; areola of membrane black (hind femora broken).

Eccritotarsus stieglmayri (Reuter, 1907)
Hsiao 1947

One female, Rio Grande do Sul, Stieglmayr. Black, head, collar, calli, basal half of hemelytra, cuneus (except apex) pale lutescent, areola and nervure of membrane, tibiae and apices of femora, antenna black, pronotum lutescent before calli.

Eurychilella pallida Reuter, 1908

One female, Bilimek, Orizaba. Cinnamon to brown, eyes, calli and posterior margin of vertex dark.

Monalocoris hesperius Reuter, 1908

One female, Bilimek, Cuerna Vacca, Oktober. Brown, very small, legs pale, second antennal segment dark apically. This species was synonymized with *eminulus* (Distant, 1893) by Carvalho, 1957: 110.

Monalocoris pallidiceps Reuter, 1907

One female, Brasilien, C. Mayr. Black; head, base of hemelytra, embolium and cuneus pale, segment II of antenna with apical half black.

Notolobus dimidiatus (Guérin-Ménéville, 1857)
Reuter, 1908

One male, Cuba, C. Mayr. Black, anterior half of body reddish (up to level of apex scutellum); median portion of disc cover in calli anteriorly.

Pachyneurhimenus pallidolimbatus Reuter, 1908

One specimen, Otto, Tombillo. Black, very similar to *Pycnoderes* Guérin-Ménéville but with