

Bothriomiris lugubris Poppius, 1915

Two specimens, Formosa, Sauter, VII, 1909 (typus, female); color varies in the two specimens (one has apex of scutellum and two subapical endocorial spots pale).

Cephalocapsidea obscurata Poppius, 1915

One specimen, Formosa, Sauter, Takan, 1897. Size minute, black, second antennal segment pale at middle, hind tibiae with minute dark spots at base of spines; head as in *Ranzovius* Distant. When studied critically this species might prove to belong to genus *Paramixia* Reuter, 1900.

Chlamydatus crassicornis Poppius, 1915

One specimen, Formosa, Sauter, Takan, 1907 (typus, male). Head above as in *Rhinacloa* Reuter, 1876, antenna pale, segment II black on apical half, femora black, tibiae with black spines but without spots at base.

Creontiades formosanus Poppius, 1915

Two specimens: (male, typus), Formosa, Sauter, Tsihorinsho; male, Formosa, Sauter, Fuhosho, 1909, VIII. Dark brown, cuneus more or less reddish, antenna pale, segment I with dark points; segment I black or black only apically.

Creontiades minutus Poppius, 1915

Four specimens: male typus, male and two females cotypes, Formosa, Sauter, Takan, 1907. Size small, pale with hind margin of pronotum and apex of scutellum dark brown, hind femur with reddish spots apically.

Creontiades pulchricornis Poppius, 1912

One female, Africa Oriental, Katona, Moschi, 1905. VIII. Pale, segment I of antenna with reddish points, frons with a red circle, sides of calli also with reddish line, propleura with red longitudinal line, abdomen red on upper margin, segment II of antenna with three reddish rings, very long.

Cylapus (Trichocylapus) clavicornis
Poppius, 1909

One specimen (typus), carded, male (?), Peru, Secuani. Scutellum cystiform, disc of pronotum, clavus and corium strongly punctate, base of cuneus pumpkin color, apex of third tibia white, second antennal segment strongly clavate, third femur with a pale fascia at middle.

Cylapus striatus Reuter, 1907

Two specimens, Peru, Pachitea (male, typus), 1 specimen Minas Gerais, Brazil, disc of pronotum with three longitudinal pale fasciae behind calli.

Deraeocoris apicalis Poppius, 1915

Two females, Formosa, Sauter. Scutellum black with basal angles and apex pale, punctate, coxae and legs pale with dark fasciae.

Under this name is also a male from Tonkin with smooth scutellum (*Camptobrochis* Fieber) and different coloration.

Deraeocoris celebensis Poppius, 1915

One male, typus and one female, S. Celebes, Bua-Kraeng, 1896, Fruhstorfer. Scutellum punctate, collar, hind margin of disc and two oblique fasciae at the sides of scutellum lutescent; size medium.

Deraeocoris colocasiae Poppius, 1915

One specimen, typus, New Guinea, Biró, 1896. Head, pronotum and legs lutescent, disc punctate, scutellum smooth, eyes large, body with erect short hairs, sternum lutescent, abdomen black.

Deraeocoris grisescens Poppius, 1915

One specimen, Australia, N.S. Wales (male, typus). Strongly punctate all over, scutellum prominent, rugose punctate, abdomen pale with a black spot at base of pygophore.