

***Deraeocoris pallidiceps* Poppius, 1915**

One male, New Guinea, Biró, 1898. Punctate (except scutellum), body with short erect hairs, brown, head and collar pale, size small.

***Deraeocoris pallidulus* Poppius, 1915**

One male, typus, Singapore, Biró, 1898. Scutellum smooth, size small, pale. The pin is strongly rusted.

***Deraeocoris papuanus* Poppius, 1915**

Two females, Erima, Austrolabe Bai, New Guinea, Biró, 1896 (typus) and New Guinea, Biró, 1897, Stephansort, Austrolabe Bai. Scutellum smooth, head and pronotum lutescent, body with short erect hairs, hemelytra dark brown.

***Deraeocoris plebejus* Poppius, 1915**

Two specimens, Formosa, Sauter, Kosempo, 1908, female (typus) and male cotypus. Light brown, cuneus reddish internally, body strongly punctate (except scutellum), embolium and corium with reddish tinge.

***Deraeocoris sauteri* Poppius, 1915**

Three specimens (female, typus), Formosa, Sauter, Fuhosho, VIII. 1908. Black, brilliant, head punctate, second antennal segment slender with long hairs, scutellum punctate, legs short, hind margin of disc narrowly pale. This species is very near if not a synonym of *Deraeocoris celebensis* Poppius.

***Eolygus vittatus* Poppius, 1915**

Two specimens (male, typus), Formosa, Sauter, Taihorinsho, 1909, VIII. Body rugose punctate with three vittae on pronotum, two on scutellum and four on hemelytra.

***Ragwelellus (Narinellus) horvathi* (Poppius, 1912) Odhiambo, 1965**

Two specimens, New Guinea, Biró, Stephansort, Australobe Bai, Biró, (typus male).

Color pumpkin, cuneus, membrane, antenna, legs (except base of femora) fuscous, eyes black. The female specimen have the femora and base of tibiae black.

***Rayeria tumidiceps* (Horvath, 1902) Odhiambo, 1965**

One female, typus, Whittom, N.S. Wales. Black, head, neck, collar, anterior area of calli at middle lutescent, exocorium with a pale sub-basal spot, abdomen with segment III (only laterally), the whole IV, VI (laterally) milky white, pygophore and segment VIII of abdomen with oblique white fasciae, body of medium size.

***Eupachypeltis flavicornis* Poppius, 1915**

One male, typus, Formosa, Sauter, Takso, 1907. Pale, size small, pubescence dense, long, erect, clavus and scutellum densely pilose, antenna with long hairs, the length larger than thickness of segments, cuneus slightly longer than wide at base.

***Eurystylomorpha crassicornis* Poppius, 1915**

One specimen, Formosa, Sauter, Fuhosho, 1908. Size medium, brown with a subapical fascia on cuneus pale, segments I and II of antenna incrossate, segments III-IV slender, the third with a pale base, clypeus compressed, prominent, seen from above.

***Eurystylus australis* Poppius, 1911**

Three specimens, Australia, Biró, 1900, N.S. Wales (typus female). As mentioned previously by Carvalho, 1976:52 this species is a synonym of *Sidnia kinbergi* (Stål, 1859).

***Eurystylus horvathi* Poppius, 1911**

One male, typus, Ceylon, Madarasz Kandy 1896, Mart.8. Size small, segment I of antenna foliaceus, membrane glassy, femora II and III pale at base, abdomen reddish with a longitudinal line of black points.