

**Eurystylus reuteri** Poppius, 1911

One specimen typus, New Guinea, Biró, 1896, Erima, Australobe Bai. The insect is lost, only label remaining.

**Felisacus longiceps** Poppius, 1915

One male, Formosa, Sauter, Chip Chip, II.1909. This species is very near if not a synonym of *magnificus* Distant, 1904, coxae, legs and abdomen pale, pygophore black, segment I of antenna incrassate towards base, apex of cuneus with reddish tinge.

**Felisacus nigricornis** Poppius, 1912

New Guinea, Biró, 1898, Sattelberg, Huon-Golf. The insect (typus) is lost.

**Fulvius brevipilis** Poppius, 1915

One specimen, New Guinea, typus.

**Fulvius brunneiceps** Poppius, 1909

Two specimens, Paraguay, Anisits, typus female; one male, Brasilia, Silvestri. Black, calli brown, second antennal segment whitish on apical third, cuneus with a large pale spot at base, corium with a transversal spot situated above apex of clavus, middle of embolium and apex of clavus pale, coxae II and III pale.

**Fulvius sauteri** Poppius, 1915

One female, typus, Formosa, Sauter, Fuhosho, VIII.1909. Apex of exocorium and embolium and apex of hind femur reddish.

**Fulvius subnitens** Poppius, 1909 and  
**Fulvius variegatus** Poppius, 1909

There are two type specimens of the first species and four specimens of the latter.

**Hallodapus persimilis** Poppius, 1915

One specimen, Formosa, Sauter, Takan, 1907. Brown, hemelytra with four pale spots (two sub-basal and two apical) on corium, two

others in the middle of clavus; cuneus and apex of hind femur reddish.

**Hyalopeplodes cyanescens** Poppius, 1912

One male, New Guinea, Biró, 1896, Lamien, Berlinhafen.

**Hyalopeplus horvathi** Poppius, 1912

One male, Africa Occidental, St. Thomé, Mocquerys.

**Isabel horvathi** Poppius, 1915

One female, Formosa, Sauter, Fuhosho, VI.1908.

**Kosmiomiris scutellaris** Poppius, 1912

One female, typus, Malacca, Perak.

**Lopidolon sordidus** Poppius, 1911

One female, Ceylon, Biró, 1902, Patypola, 2000 m. This species belongs to genus *Bromeliaemiris* Schumacher, 1919 (*n. comb.*). Color brown, vertex and margin of eyes, longitudinal fasciae on middle of pronotum and collar, scutellum, embolium and cuneus (except black apex of the latter) lutescent; calli, eyes, antenna and frons black, a longitudinal fascia along propleura lutescent.

**Lygus bakeri** Poppius, 1915

Four specimens, Formosa, Sauter, Takan. Color and aspect of *Taylorilygus pallidulus* (Blanchard, 1852), apex of hind femur black with two reddish rings antennal segments III-IV dark, tibiae with small dark points at base of spines; in some specimens apex of clavus and apex of embolium black, size small.

**Lygus basalis** Poppius, 1914

Two specimens (male and female), Singapore, Biró, 1898. Small, eyes large, straw color, two small spots at apex of embolium and cuneus black, hind femur with two subapical reddish rings, tibiae with dark points at base of spines,