

fasciae on pronotum, scutellum reddish (except basal angles and apex which are pale), apex of cuneus black.

***Lygus punctiscutum* Poppius, 1914**

One female, typus, Singapore, Biró, 1898. Body strongly punctate, including scutellum which is very prominent, glabrous, head and scutellum whitish, clavus (except base and apex) and apex of embolium black, two small spots near humeral angles, hind margin of disc white; size medium, legs and clypeus pumpkin color. It belongs probably to genus *Deraeocoris* Kirschbaum.

***Lygus rufescens* Poppius, 1914**

One female, Borneo, Brunei. Testaceous, underside of body reddish, cuneus red; size small. This species is a homonym of *Lygus rufescens* Hahn, 1831 (Wanz.Ins.1.28, fig. 18) which was synonymized by Reuter, 1878 (Hem. Gymn. Eur.1:138) with *Psallus varians* Herrich-Schaeffer, 1841. Therefore it is proposed here to rename it *Lygus bruneiensis* nom. nov.

***Lygus sauteri* Poppius, 1912**

Four specimens (typus, male), Formosa, Sauter, Ching Ching, 1909, II. Abdomen, meso and metasternum black, xyphus lutescent, size small, brilliant, scutellum white with a dark spot at middle of base, corium and embolium with pale spot externally, apex of corium and subapical portion of cuneus pale yellow.

***Lygus stali* Poppius, 1914**

One male, typus, New Guinea, Biró, 1896. Straw color, apex of second antennal segment, segments III-IV black, size small.

***Lygus sublaevis* Poppius, 1914**

One female, typus, S. Celebes, Bua-Kraeng, H. Fruhstorfer, 5000, Febr., 1896. Small, straw color, cuneal fracture deep and wide, cuneus with a brown spot at middle, segment I of antenna at base and apex, segment II and III at apices black, membrane translucent, larger areola rounded.

***Lygus taivanensis* Poppius, 1915**

Two specimens, female, typus, Formosa, Sauter, Taiwan, 1909.II. Small, color greenish straw, a spot on clavus near apex of scutellum, apex of clavus, a transversal irregular fascia on apex of hemelytra and apex of cuneus black; femur III marmorate with brown, tibiae III with dark spots at base of spines; antennae III and IV black.

***Lygus taivanus* Poppius, 1915**

One female, typus (four cotypes), Formosa, Sauter, Chip Chip, 1909, II. Large, elongate, color straw tending to greenish, femur III marmorate with red, cuneus very long, about three times as long as wide at base, apex of segment II plus segments III and IV black, tarsi III black. This species does not belong to genus *Lygus* Hahn and might be included in the future in the genus *Calocoris* Fieber.

***Lygus umbratus* Poppius, 1914**

One female, typus, Samanga, South Celebes, H. Fruhstorfer, Nov. 1895. Size small, color straw, tibiae III red at base and with black spots at base of spines, apex of clavus, apex of embolium and apex of cuneus black, femora III with two reddish rings. This species is very near if not a synonym of *Lygus bakeri* Poppius, 1915.

***Lygus viridiflavus* Poppius, 1914**

One female, typus, Darjeeling, Fruhstorfer. Medium size, color straw, translucent, tending to lutescent or citrine, antenna III and IV black, rostrum reaching second coxae. This species might belong to genus *Pleurochilophorus* Reuter, 1905.

***Lygus vittulatus* Poppius, 1914**

One female, typus, South Celebes, Bua-Kraeng, H. Fruhstorfer, 5000 m, Feb., 1895. Size large, color straw, hind margin of disc (a transversal sub-basal spot), a longitudinal spot on scutellum (except apex), claval and corio-claval commissures, external margin and subapical spot of embolium, apex of cuneus dark. The